

78505 - 506.3 grams

78506 - 56 grams

78507 - 23.4 grams

78509 - 8.7 grams

Ilmenite Basalt



Figure 1: Photo of 78505. Scale and cube are in cm. S73-15386. White line shows approximate position of saw cut.




Figure 2: Map of station 8 showing location of soil 78500 and rake sample 78525-78599.




Figure 3: Composition of pyroxene and olivine in 78505 (Warner et al. 1978).




Figure 4: Photo of 78506. Scale in cm. S73-15468.

Mineralogical Mode

	Warner	Brown
Olivine	0.5	0.5
Pyroxene	45.7	47.7
Plagioclase	35.7	27.7
Opakes	15.3	21
Silica	1.5	1.9
Meostasis	1.1	1.2

Introduction

According to Warner et al. (1978) 78505 is a plagioclase-poikilitic ilmenite basalt similar to 71565, 71567 and 71509. It has a coarse grain size (1 mm) with abundant pigeonite and blocky ilmenite (figures 6 and 7).

78505 – 78509 were collected as part of a comprehensive soil sample (78500) at station 8, Apollo 17. A large rake sample collected adjacent to 78500 also returned abundant mare basalt fragments.

Petrography

Brown et al. (1975) and others studied the Apollo 17 mare basalts including 78505 finding that they are all ilmenite-rich because of their high Ti content. 78505 is slightly different because of its proportion of low-Ca pigeonite (figure 3).



Figure 5: Photo of 78509. About 2 cm across. S73-18608

Table 2: Zr-rich mineral 78506.

(Brown et al. 1975)

TiO ₂	28.8
FeO	9.71
MgO	0.22
CaO	3.15
ZrO ₂	35.13
Y ₂ O ₃	8.31
NbO ₅	2.7

Warner et al. (1978) reported tranquillityite and zirconolite in 78505. Brown et al. (1975) reported a Zr-rich mineral in 78506 (table 2). Roedder (1979) discusses the low-K inclusions in ilmenite in 78505.

Usselman et al. (1975) and others experimentally reproduced the textures and mineral chemistries of high-titanium mare basalts from Apollo 17 and discussed cooling rates.

Chemistry

The chemical compositions of 78505, 78506, 78507 and 78509 are tabulated in table 1. These samples are typical of other Apollo 17 mare basalts (figures 10, 11 and 12).

Cosmogenic isotopes and exposure ages

Drozd et al. (1977) determined the cosmic ray exposure age to be 121 m.y.

Keith et al. (1974) reported the cosmic-ray-induced activity of 78505.


Other Studies

Stephenson et al. (1975) reported studies of the magnetic properties of 78505.

Processing

78505 was sawn in half (figure 8) but it broke in pieces. There are 12 thin sections of 78505, 3 for 78506 and one each for 78507 and 78509.

*Figure 6: Texture of 78505,61.
Field of view is 2.8 mm across.*






Lunar Sample Compendium
C Meyer 2011



*Figure 7: Photo of thin section 78506,27
with crossed Nicols. 2.8 mm across*







Figure 8: Processing photo for 78505. S74-19019 Cube is 1 cm³.






Figure 9: Magnetic properties of 78505 (Stephenson et al. 1975).




Figure 10: Composition of 78505 compared with other Apollo basalts.




Figure 11: Normalized rare-earth-element diagram for 78505 compared with A and B types of Apollo 17 basalt.




Figure 12: Trace element ratios for 78506 compared with that of other Apollo 17 basalts.

Table 1. Chemical composition of 78505.

	78505		78506		78507		78509	
<i>reference weight</i>	Warner75	Keith 74	Rhodes76	Nyquist76	Warner79	Warner79		
SiO ₂ %			38.55	(d)				
TiO ₂	12	(a)	12.93	(d)	11.9	12.3	(a)	
Al ₂ O ₃	10.6	(a)	8.99	(d)	8.8	9.2	(a)	
FeO	18.6	(a)	19.36	(d)	18	19	(a)	
MnO	0.227	(a)	0.27	(d)	0.222	0.252	(a)	
MgO	9.5	(a)	9.59	(d)	10	8	(a)	
CaO	9.9	(a)	9.94	(d)	9.7	10.9	(a)	
Na ₂ O	0.458	(a)	0.39	(d)	0.407	0.414	(a)	
K ₂ O	0.07	(a)	0.061	(b)	0.05	(d)	0.037	0.04 (a)
P ₂ O ₅			0.02	(d)				
S %			0.16	(d)				
<i>sum</i>								
Sc ppm	74	(a)	73	(b)	79	89	(a)	
V	108	(a)			130	101	(a)	
Cr	2983	(a)	3489	(d)	3856	2657	(a)	
Co	18.7	(a)	17.6	(b)	21	22	(a)	
Ni								
Cu								
Zn								
Ga								
Ge ppb								
As								
Se								
Rb			0.44	(c)				
Sr			175	(c)				
Y								
Zr								
Nb								
Mo								
Ru								
Rh								
Pd ppb								
Ag ppb								
Cd ppb								
In ppb								
Sn ppb								
Sb ppb								
Te ppb								
Cs ppm								
Ba			66	(c)				
La	5.9	(a)	5.1	(c)	3.4	3.9	(a)	
Ce			17.8	(c)	13	14	(a)	
Pr								
Nd			19.6	(c)	16	16	(a)	
Sm	9.4	(a)	8.19	(c)	6	5.8	(a)	
Eu	2.1	(a)	1.85	(c)	1.59	1.22	(a)	
Gd			12.9	(c)				
Tb					1.4	1.3	(a)	
Dy	15	(a)	14.9	(c)	10	9	(a)	
Ho								
Er								
Tm								
Yb	8.9	(a)	7.99	(c)	5.8	5.5	(a)	
Lu	1.2	(a)	1.11	(b)	0.86	0.8	(a)	
Hf			8.2	(b)	5.5	5.1	(a)	
Ta					1.3	1.3	(a)	
W ppb								
Re ppb								
Os ppb								
Ir ppb								
Pt ppb								
Au ppb								
Th ppm		0.39	(b)					
U ppm		0.135	(b)					

technique: (a) INAA, (b) radiation count. (c) IDMS, (d) XRF

References for 78505, 78506, 78507 and 78509.

- Brown G.M., Peckett A., Emeleus C.H., Phillips R. and Pinsent R.H. (1975a) Petrology and mineralogy of Apollo 17 mare basalts. *Proc. 6th Lunar Sci. Conf.* 1-13.
- Butler P. (1973) **Lunar Sample Information Catalog Apollo 17.** Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.
- Drozd R.J., Hohenberg C.M., Morgan C.J., Podosek F.A. and Wroge M.L. (1977) Cosmic-ray exposure history at Taurus-Littrow. *Proc. 8th Lunar Sci. Conf.* 3027-3043.
- Gibson E.K., Usselman T.M. and Morris R.V. (1976a) Sulfur in the Apollo 17 basalts and their source regions. *Proc. 7th Lunar Sci. Conf.* 1491-1505.
- Keith J.E., Clark R.S. and Bennett L.J. (1974a) Determination of natural and cosmic ray induced radionuclides in Apollo 17 lunar samples. *Proc. 5th Lunar Sci. Conf.* 2121-2138.
- LSPET (1973) Apollo 17 lunar samples: Chemical and petrographic description. *Science* **182**, 659-672.
- LSPET (1973) Preliminary Examination of lunar samples. Apollo 17 Preliminary Science Rpt. NASA SP-330. 7-1 – 7-46.
- Ma M.-S., Schmitt R.A., Warner R.D., Taylor G.J. and Keil K. (1979b) Composition, petrography, and genesis of Apollo 17 high-Ti mare basalts (abs). *Lunar Planet. Sci.* **X**, 765-767. Lunar Planetary Institute, Houston.
- Muehlberger W.R. and many others (1973) Preliminary Geological Investigation of the Apollo 17 Landing Site. *In Apollo 17 Preliminary Science Report.* NASA SP-330.
- Neal C.R. and Taylor L.A. (1993) Catalog of Apollo 17 rocks. Vol. 3 Central Valley
- Nyquist L.E., Bansal B.M. and Wiesmann H. (1976a) Sr isotopic constraints on the petrogenesis of Apollo 17 mare basalts. *Proc. 7th Lunar Sci. Conf.* 1507-1528.
- Roedder E. (1979a) Melt inclusions in 75075 and 78505 - the problem of anomalous low-K inclusions in ilmenite revisited. *Proc. 10th Lunar Planet. Sci. Conf.* 249-257.
- Rhodes J.M., Hubbard N.J., Wiesmann H., Rodgers K.V., Brannon J.C. and Bansal B.M. (1976a) Chemistry, classification, and petrogenesis of Apollo 17 mare basalts. *Proc. 7th Lunar Sci. Conf.* 1467-1489.
- Stephenson A., Runcorn S.K. and Collinson D.W. (1975) On changes in intensity of the ancient lunar magnetic field. *Proc. 6th Lunar Sci. Conf.* 3049-3062.
- Usselman T.M., Lofgren G.E., Donaldson C.H. and Williams R.J. (1975) Experimentally reproduced textures and mineral chemistries of high-titanium mare basalts. *Proc. 6th Lunar Sci. Conf.* 997-1020.
- Warner R.D., Keil K., Murali A.V. and Schmitt R.A. (1975a) Petrogenetic relationships among Apollo-17 basalts. *In Papers presented to the Conference on Origins of Mare Basalts and their Implications for Lunar Evolution* (Lunar Science Institute, Houston), 179-183.
- Warner R.D., Keil K., Nehru C.E. and Taylor G.J. (1978) Catalogue of Apollo 17 rake samples from Stations 1a, 2, 7, and 8. Spec. Publ. #18, UNM Institute of Meteoritics, Albuquerque. 88 pp.
- Warner R.D., Nehru C.E. and Keil K. (1978g) Opaque oxide mineral crystallization in lunar high-titanium basalts. *Am. Mineral.* **68**, 1209-1224.
- Warner R.D., Taylor G.J., Conrad G.H., Northrop H.R., Barker S., Keil K., Ma M.-S. and Schmitt R. (1979) Apollo 17 high-Ti mare basalts: New bulk compositional data, magma types, and petrogenesis. *Proc. 10th Lunar Planet. Sci. Conf.* 225-247.
- Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L. and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.