

10058

Ilmenite Basalt (low K)
282 grams



Figure 1: Photo of 10058 after breaking. Large piece is 5 cm. NASA S69-47475.

Introduction

Lunar sample 10058 is a low-K, ilmenite basalt with a relatively coarse texture (figure 1). It has a crystallization age of 3.7 b.y. with a cosmic ray exposure age of 70 m.y.

Petrography

Schmitt et al. (1970) termed 10058 as a “medium-grained, vuggy, ophitic, cristobalite basalt.” James and Jackson (1970) termed it a medium grained ophitic basalt. Brown et al. (1970) termed the texture of 10058 as “gabboic”, but found no evidence for crystal accumulation or flow orientation. Beaty and Albee (1978) note that 10044, 10047 and 10058 “are so similar to one another that it seems quite likely that these rocks are fragments of a larger block.” Ilmenite grains up to 2 mm are reported (Simpson and Bowie 1970).

Brown et al. (1970) and Beaty and Albee (1978) studied the mineral chemistry in 10058.

Mineralogy

Olivine: Beaty and Albee (1978) found one rounded, anhedral olivine grain mantled by a large zoned pyroxene. The olivine is chemically zoned (Fo_{68-58}). Brown et al. (1970) reported trace fayalite in the mesostasis.

Pyroxene: Hargraves et al. (1970) and Beaty and Albee (1978) studied the sector-zoned pyroxene in 10058. Brown et al. reported optically-visible pigeonite exsolution.

Amphibole (?): Gay et al. (1970) reported finding dark green clinoamphibole in a vug.




Figure 2: Photomicrographs of thin section of 10058. Top is plane-polarized light and bottom is crossed Nicols. Scale is 2.5 mm. Note patch of silica in center shaped like a "chinese character". NASA S70-49011 and 12.

Plagioclase: Beaty and Albee (1978) found plagioclase was An_{94-70} , while Brown et al. (1970) reported An_{92-82} .

Ilmenite: Simpson and Bowie (1970) studied the complex intergrowth of ilmenite with ulvöspinel in 10058 and give an analysis of ilmenite.

Y, Zr silicate: Cameron (1970) reported a new mineral – probably tranquillityite.

Troilite: Simpson and Bowie (1970) give an analysis.

Phosphate: Brown et al. reported trace apatite.

Cristobalite: Silica in 10058 has a distinctive cracking pattern, due to inversion of high to low cristobalite structure (Nord et al. 1974).




Figure 3: Pyroxene and olivine composition of 10058 (from Beatty and Albee 1978).

Chemistry

The chemical composition of 10058 is tabulated in table 1 and figures 4 and 5.

Radiogenic age dating

Papanastassiou et al. (1970) determined the Rb/Sr isochron age for 10058. Guggisberg et al. (1979) determined the age by the Ar/Ar plateau technique (figure 6).

Cosmogenic isotopes and exposure ages

Guggisberg et al. (1979) determined an $^{37}\text{Ar}/^{38}\text{Ar}$ exposure age of 71 m.y. Eberhardt et al. (1970) calculated an exposure age of 65 m.y. from the Ar data by Funkhouser et al. (1970) and Pepin et al. (1970).

Other Studies

Oxygen isotopes were reported for mineral separates of 10058 by Onuma et al. (1970) and Taylor and Epstein (1970).

Mineralogical Mode	Beatty and Albee 1978	Brown et al. 1970
Olivine		
Pyroxene	47	45.7
Plagioclase	34	37.1
Ilmenite	13	10.5
Glass	0.05	
silica	4.8	5.1
troilite	0.43	.27
phosphate	0.42	




Figure 4: Composition of 10058 compared with that of other Apollo lunar samples.

Pepin et al. (1970), Funkouser et al. (1970) and Bogard et al. (1971) reported the abundance and isotopic composition of rare gasses from 10058.

Crozaz et al. (1970) studied the cosmic ray tracks as a function of depth in plagioclase crystals for 10058 (figure 7) and determined the “erosion rate” by micrometeorite bombardment by comparing the apparent exposure age by tracks with the exposure age determined by ^{81}Kr and ^{38}Ar .

Housley et al. (1970) determined the Mossbauer spectra.

Processing

Apollo 11 samples were originally described and cataloged in 1969 and “re-cataloged” by Kramer et al. (1977). There are 12 thin sections.

List of Photo #s for 10058

S69-46297 – 320	B&W mug
S69-47473 – 6	
S69-47485 – 6	
S70-49009 – 12	TS color
S70-49967 – 8	TS
S70-49874 – 5	TS
S74-27032	,34
S76-21354 – 5	
S76-26326 – 7	TS
S76-23295 – 6	,109




Figure 5: Normalized rare-earth-element composition for low-K basalt 10058 (the line) compared with that of low-K basalt 10020 and high-K basalt 10049 (the dots) (data from Wiesmann et al. 1975).




Figure 6: Argon release pattern for 10044 and 10058 (Guggisberg et al. 1979).

Summary of Age Data for 10058

Rb/Sr	Ar/Ar plateau
Papanastassiou et al. 1970	3.63 ± 0.20 b.y.
Guggisberg et al. 1979	3.71 ± 0.04
Caution: Change in Rb decay constant.	

Table 1. Chemical composition of 10058.

reference weight	LSPET69	Tera70	Dickinson89	Rose70	Gast70	Wiesmann75	Goles70	Rhodes80
SiO ₂ %	43	(a)		41.4	41.7	(d)	40.6	(c) 42.18 (e) 1.51 g
TiO ₂	9	(a)		11.1	9.55	(d)	10.2	(c) 9.37 (e)
Al ₂ O ₃	13	(a)		10.7	11.8	(d)	11.1	(c) 10.45 (e)
FeO	17	(a)	15.2	(c) 17.3	18.2	(d)	19.2	(c) 18.96 (e)
MnO	0.55	(a)		0.27	0.27	(d)	0.24	(c) 0.3 (e)
MgO	6.5	(a)		6.25	6.3	(d)	6.3	(c) 5.91 (e)
CaO	10.5	(a) 12	(b) 8.4	(c) 12.1	11	(d)	11.3	(c) 12.28 (e)
Na ₂ O	0.56	(a) 0.44	0.4	0.79	0.68	(d) 0.39	39	0.41 (c) 0.39 (c)
K ₂ O	0.11	(a) 0.1	(b)	0.07	0.09	(d) 0.1	(b) 0.1	0.11 (e) 0.13 (e)
P ₂ O ₅								
S %								
sum								
Sc ppm			76	(c)			80.8	(c) 88 (c)
V	32	(a)					78	(c)
Cr	3700	(a)	1400	(c) 1437	1642	(d)	1800	(c) 1390 (c)
Co	7	(a)	10	(c)			14.4	(c) 11 (c)
Ni								
Cu								
Zn								
Ga			12	(c)			8.4	(e)
Ge ppb			2.7	(c)				
As								
Se								
Rb	1.6	(a) 1.02	(b)			0.98	(b) 0.98	(b) 1.2 (e)
Sr	190	(a)	186	(c)		218	(b) 218	(b) 224 (e)
Y	230	(a)						141 (e)
Zr	250	(a)					190	(c)
Nb								
Mo								
Ru								
Rh								
Pd ppb								
Ag ppb								
Cd ppb								
In ppb								
Sn ppb								
Sb ppb								
Te ppb								
Cs ppm		0.04	(b)			0.03	(b) 0.027	(b)
Ba	85	(a) 113	(b) 122	(c)		117	(b) 117	(b) 140 (c)
La			9.7	(c)		11.5	(b) 11.5	(b) 11.8 (c) 10.3 (c)
Ce			33	(c)		40.2	(b) 40.2	(b) 39 (c) 40 (c)
Pr								
Nd			20	(c)		41.2	(b) 41.2	(b)
Sm			41	(c)		17.2	(b) 17.2	(b) 14 (c) 16.4 (c)
Eu			2.3	(c)		2.64	(b) 2.64	(b) 2.14 (c) 2.42 (c)
Gd						23.6	(b) 23.6	(b)
Tb			4.3	(c)				3.5 (c) 3.4 (c)
Dy						27	(b) 27	(b)
Ho								5.5 (c)
Er			0.57	(c)		16.3	(b) 16.3	(b)
Tm			0.71	(c)				
Yb	5	(a)	13.5	(c)		15.5	(b) 13.8	14 (c) 13.3 (c)
Lu			2.1	(c)		2.14	(b) 2.14	(b) 1.94 (c) 1.9 (c)
Hf			10.8	(c)				11.2 (c) 12.1 (c)
Ta			1.9	(c)				1.6 (c) 2.5 (c)
W ppb								
Re ppb								
Os ppb								
Ir ppb								
Pt ppb								
Au ppb								
Th ppm			0.84	(c)				1.2 (c)
U ppm							0.18 (c)	

technique: (a) OES, (b) IDMS, (c) INAA, (d) semimicro XRF, (e) XRF

Table 1b. Chemical composition of 10058.

reference	Duncan76	Morrison70	Beaty78	Murthy70	Hurley70	Neal2001
<i>weight</i>						
SiO ₂ %	41.78	(e) 39.4	(f) 40.24	39.76	(g)	
TiO ₂	9.07	(e) 9.7	(f) 10.6	10.52	(g)	
Al ₂ O ₃	10.44	(e) 10.2	(f) 10.34	10.54	(g)	
FeO	18.68	(e) 19.6	(f) 19.6	19.36	(g)	
MnO	0.266	(e) 0.25	(f) 0.29	0.28	(g)	
MgO	5.96	(e) 5.6	(f) 5.8	6.22	(g)	
CaO	12.34	(e) 15.4	(f) 12.18	12.53	(g)	
Na ₂ O	0.48	(e) 0.43	(f) 0.42	0.42	(g)	
K ₂ O	0.085	(e) 0.11	(f) 0.06	0.02	(g) 0.098	(b)
P ₂ O ₅	0.132	(e) 0.055	(f) 0.16	0.09	(g)	
S %	0.19	(e)		0.2	0.22	(g)
sum	99.423					
Sc ppm		87	(f)		88.6	(h)
V	46	(e) 41	(f)		65.3	(h)
Cr	1402	(e) 1500	(f)		1201	(h)
Co	12	(e) 14	(f)		15.5	(h)
Ni	<2	(e)			1.15	(h)
Cu		7.1	(f)		37.6	(h)
Zn		9.3	(f)		88	(h)
Ga		4.3	(f)		4.5	(h)
Ge ppb						
As		70	(f)			
Se						
Rb	1.5	(e) 1.2	(f)	1.15	(b) 0.68	(b) 0.88
Sr	208	(e) 180	(f)	224	(b) 172	(b) 217
Y	147	(e) 150	(f)			111
Zr	376	(e) 380	(f)			300
Nb	28.4	(e) 47	(f)			25
Mo		0.4	(f)			0.19
Ru						
Rh						
Pd ppb		200	(f)			
Ag ppb		70	(f)			
Cd ppb		700	(f)			
In ppb		600	(f)			
Sn ppb		1200	(f)			
Sb ppb		10	(f)			
Te ppb						
Cs ppm		0.3	(f)		0.05	(h)
Ba	136	(e) 140	(f)	128	(b)	108
La		16	(f)			11.5
Ce		45	(f)			45.4
Pr		13	(f)			7.33
Nd		72	(f)			42.9
Sm		22	(f)			18.2
Eu		3	(f)			2.4
Gd		22	(f)			23.3
Tb		5.4	(f)			4.5
Dy		39	(f)			28.1
Ho		9	(f)			5.56
Er		36	(f)			16.1
Tm		2	(f)			2.25
Yb		22	(f)			15.3
Lu		2.3	(f)			2.07
Hf		13	(f)			11.2
Ta		1	(f)			1.82
W ppb		360	(f)			160
Re ppb						
Os ppb						
Ir ppb						
Pt ppb						
Au ppb						
Th ppm		1.1	(f)		0.74	(h)
U ppm		0.2	(f)		0.23	(h)

technique: (e) XRF, (f) various, (g) elec. Probe, (h) ICP-MS





Figure 7: Etched cosmic ray track profile with depth for plagioclase (circles) from 10058 (Crozaz et al. 1970).

References for 10058

- Beaty D.W. and Albee A.L. (1978) Comparative petrology and possible genetic relations among the Apollo 11 basalts. *Proc. 9th Lunar Planet. Sci. Conf.* 359-463.
- Bogard D.D., Funkhouser J.G., Schaeffer O.A. and Zahringer J. (1971) Noble gas abundances in lunar material-cosmic ray spallation products and radiation ages from the Sea of Tranquillity and the Ocean of Storms. *J. Geophys. Res.* **76**, 2757-2779.
- Brown G.M., Emeleus C.H., Holland J.G. and Phillips R. (1970) Mineralogical, chemical, and petrological features of Apollo 11 rocks and their relationship to igneous processes. *Proc. Apollo 11 Lunar Sci. Conf.* 195-219.
- Cameron E.N. (1970) Opaque minerals in certain lunar rocks from Apollo 11. *Proc. Apollo 11 Lunar Sci. Conf.* 221-245.
- Crozaz G., Haack U., Hair M., Maurette M., Walker R. and Woolum D. (1970) Nuclear track studies of ancient solar radiations and dynamic lunar surface processes. *Proc. Apollo 11 Lunar Science Conf.* 2051-2080.
- Duncan A.R., Erlank A.J., Willis J.P., Sher M.K. and Ahrens L.H. (1974a) Trace element evidence for a two-stage origin of some titaniferous mare basalts. *Proc. 5th Lunar Sci. Conf.* 1147-1157.
- Dickinson T., Taylor G.J., Keil K. and Bild R.W. (1989) Germanium abundances in lunar basalts: Evidence of mantle metasomatism. *Proc. 19th Lunar Planet. Sci.* 189-198. Lunar Planetary Institute, Houston.
- Eberhardt P., Geiss J., Graf H., Grogler N., Krahenbuhl U., Schwaller H., Schwarzmuller J. and Stettler A. (1970) Correlation between rock type and irradiation history of Apollo 11 igneous rocks. *Earth Planet. Sci. Lett.* **10**, 67-72.
- Funkhouser J.G., Schaeffer O.A., Bogard D.D. and Zahringer J. (1970) Gas analysis of the lunar surface. *Proc. Apollo 11 Lunar Sci. Conf.* 1111-1116.
- Gast P.W., Hubbard N.J. and Wiesmann H. (1970b) Chemical composition and petrogenesis of basalts from Tranquillity Base. *Proc. Apollo 11 Lunar Sci. Conf.* 1143-1163.
- Gay P., Bancroft G.M. and Brown M.G. (1970) Diffraction and Mossbauer studies of minerals from lunar soils and rocks. *Proc. Apollo 11 Lunar Sci. Conf.* 481-497.
- Goles G.G., Randle K., Osawa M., Lindstrom D.J., Jerome D.Y., Steinborn T.L., Beyer R.L., Martin M.R. and McKay S.M. (1970) Interpretations and speculations on elemental abundances in lunar samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1177-1194.
- Guggisberg S., Eberhardt P., Geiss J., Grogler N., Stettler A., Brown G.M. and Pecket A. (1979) Classification of the Apollo-11 basalts according to Ar³⁹-Ar⁴⁰ ages and petrological properties. *Proc. 10th Lunar Planet. Sci. Conf.* 1-39.
- Hollister L.S. and Hargraves R.B. (1970) Compositional zoning and its significance in pyroxenes from two coarse grained Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 541-550.
- Housley R.M., Blander M., Abdel-Gawad M., Grant R.W. and Muir A.H. (1970) Mossbauer spectroscopy of Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 2251-2268.
- Hurley P.M. and Pinson W.H. (1971) Whole-rock Rb-Sr isotopic age relationships in Apollo 11 lunar samples. *Proc. Apollo 11 Lunar Sample Conference* 1311-1315.
- James O.B. and Jackson E.D. (1970) Petrology of the Apollo 11 ilmenite basalts. *J. Geophys. Res.* **75**, 5793-5824.
- Kramer F.E., Twedell D.B. and Walton W.J.A. (1977) **Apollo 11 Lunar Sample Information Catalogue** (revised). Curator's Office, JSC 12522
- LSPET (1969) Preliminary examination of lunar samples from Apollo 11. *Science* **165**, 1211-1227.
- Morrison G.H., Gerard J.T., Kashuba A.T., Gangadharam E.V., Rothenberg A.M., Potter N.M. and Miller G.B. (1970) Elemental abundances of lunar soil and rocks. *Proc. Apollo 11 Lunar Sci. Conf.* 1383-1392.
- Murthy V.R., Evensen N.M., Jahn B.-M. and Coscio M.R. (1971) Rb-Sr ages and elemental abundances of K, Rb, Sr and Ba in samples from the Ocean of Storms. *Geochim. Cosmochim. Acta* **35**, 1139-1153.

- Neal C.R. (2001) Interior of the moon: The presence of garnet in the primitive deep lunar mantle. *J. Geophys. Res.* **106**, 27865-27885.
- Nord G.L., Lally J.S., Heuer A.H., Christie J.M., Radcliffe S.V., Fisher R.M. and Griggs D.T. (1974) A mineralogical study of rock 70017, an ilmenite-rich basalt, by high voltage electron microscopy (abs). *Lunar Sci. V*, 556-558. Lunar Planetary Institute, Houston
- Onuma N., Clayton R.N. and Mayeda T. (1970) Apollo 11 rocks: Oxygen isotope fractionation between minerals and an estimate of the temperature of formation. *Proc. Apollo 11 Lunar Sci. Conf.* 1429-1434.
- Papanastassiou D.A., Wasserburg G.J. and Burnett D.S. (1970a) Rb-Sr ages of lunar rocks from the Sea of Tranquillity. *Earth Planet. Sci. Lett.* **8**, 1
- Pepin R.O., Nyquist L.E., Phinney D. and Black D.C. (1970) Rare gases in Apollo 11 lunar material. *Proc. Apollo 11 Lunar Sci. Conf.* 1435-1454.
- Rhodes J.M. and Blanchard D.P. (1980) Chemistry of Apollo 11 low-K mare basalts. *Proc. 11th Lunar Planet. Sci. Conf.* 49-66.
- Rose H.J., Cuttitta F., Dwornik E.J., Carron M.K., Christian R.P., Lindsay J.R., Ligon D.T. and Larson R.R. (1970b) Semimicro X-ray fluorescence analysis of lunar samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1493-1497.
- Schmitt H.H., Lofgren G., Swann G.A. and Simmons G. (1970) The Apollo 11 samples: Introduction. *Proc. Apollo 11 Lunar Science Conf.* 1-54.
- Simpson P.R. and Bowie S.H.U. (1970) Quantitative optical and electron-probe studies of opaque phases in Apollo 11 samples. *Proc. Apollo 11 Lunar Sci. Conf.* 873-890.
- Sutton R.L. and Schaber G.G. (1971) Lunar locations and orientations of rock samples from Apollo missions 11 and 12. *Proc. 2nd Lunar Sci. Conf.* 17-26.
- Taylor H.P. and Epstein S. (1970a) O18/O16 ratios of Apollo 11 lunar rocks and minerals. *Proc. Apollo 11 Lunar Sci. Conf.* 1613-1626.
- Tera F., Eugster O., Burnett D.S. and Wasserburg G.J. (1970) Comparative study of Li, Na, K, Rb, Cs, Ca, Sr, and Ba abundances in achondrites and in Apollo 11 lunar samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1637-1657.
- Turner G. (1970a) Argon-40/argon-39 dating of lunar rock samples. *Proc. Apollo 11 Lunar Sci. Conf.* 1665-1684.
- Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample Data Generated by the Gast, Nyquist and Hubbard Lunar Sample PI-Ships. Unpublished. JSC