

12035
Olivine Basalt
71 grams




Figure 1: Photo of 12035. NASA #S69-61249.

Introduction

Lunar basalt 12035 was found on the rim of Bench Crater. It is a friable, vuggy, micro-gabbro with mafic composition that initially broke into 6+ smaller pieces during PET. Figures 1, 2 and 4 show the hackly, coarse grain, nature of 12035. It was mistakenly termed “troctolite” in the Apollo 12 catalog (Warner 1971).

Petrography

James and Wright (1972) describe 12035 as an “olivine gabbro” with euhedral to subhedral chrome spinel and olivine, and subhedral to anhedral clinopyroxene enclosed by poikilitic plagioclase. Interstitial ilmenite tends to be anhedral. Reid (1971) also described 12035 as a “coarse olivine-rich (30%) gabbro with euhedral clinopyroxene, subhedral olivine in large plagioclase

Mineralogical Mode for 12035

	Neal et al. 1994	Brett et al. 1971	Papike et al. 1976	Walker et al. 1976
Olivine	21.4	36	32	28.7
Pyroxene	51.1	32.6	41.6	
Plagioclase	24.3	27.2	19.7	
Opaques			6.2	
Ilmenite	0.1	2.5		
Chromite +Usp	2.3	1		
mesostasis	0.6	0.6	0.5	
“silica”				



Figure 2: Photo of 12035,9 with mm scale. NASA ↑
S70-44176.

grains". The olivine is relatively iron rich, hence it is not a simple cumulate for the other olivine basalts, but rather the olivine has re-equilibrated with the liquid (Butler 1976).

Walker et al. (1976) studied the sequence of olivine basalts (figure 8) and place 12035 about 11 meter above the base of the flow in their model.

Olivine phenocrysts in 12035 contains melt inclusions (see figure 1-11 in Roedder and Weiblen 1971).

Mineralogy

Olivine: Olivine in 12035 is Fe-rich (Fo_{64}) although the first olivine to crystallize should have been Mg-rich (Fo_{80+}), indicating that olivine in 12035 re-equilibrated sub-solidus. Butler (1973) determined the minor element content of olivine (figure 6).

Pyroxene: Schnetzler and Phillipotts (1971) studied the distribution of trace elements between pyroxene and plagioclase in 12035.

Plagioclase: Walker et al. (1976) found that 12035 had the largest grains of plagioclase (~1 mm) of the Apollo 12 olivine basalts.

↓ Figure 3: Photomicrographs of 12035,23. Scale is
2.6 mm. NASA # S70-49465-466.







Figure 4: Photo of the other side of 12035,9. NASA photo # S70-44178.




Figure 5: Photo of thin section 12035,21. 1.5 cm across. NASA #S70-45637.

Spinel: The spinel in 12035 has been carefully studied by Reid (1971) who showed that the spinel re-equilibrated.




Figure 6: Pyroxene and olivine composition of 12035.

Metal: Brett et al. (1971) determined the Ni content of minute metallic iron grains in 12035 (figure 7).

Chemistry

Compston et al. (1971) reported the major and some minor element composition of 12035 (figure 10). Trace




Figure 7: Histogram of Ni concentrations of metal grains in 12035 and 12040 lunar samples (lifted from Brett et al. 1971).




Figure 8: Size of plagioclase grains compared with normative amount of olivine. (Walker et al. 1976).

elements were determined by Gast and Hubbard (1970), Cuttitta et al. (1971) and Hubbard et al (1971) (figure 9).

Radiogenic age dating

Papanastassiou and Wasserburg (1971a) reported a Rb/Sr age of 3.2 b.y. for 12035 (figure 11).

Cosmogenic isotopes and exposure ages

Burnett et al. (1975) determined an exposure age of 115 ± 45 m.y. by $^{81}\text{Kr}/^{83}\text{Kr}$.

Processing

12035 initially broke into 6 pieces plus fines. Most early allocations were from 12036,9; recent allocations were from 12035,1. There are 9 thin sections.




Figure 9: Normalized rare-earth-element diagram for 12035 (data from Hubbard and Gast 1971 and Wiesmann et al. 1975).




Figure 10: Composition of 12035 compared with other lunar basalts.




Figure 11: Rb-Sr isochron for 12035 (Papanastassiou and Wasserburg 1971a).


Summary of Age Data for 12035

Rb/Sr
Papanastassiou and Wasserburg 1971a 3.2 b.y

Table 1. Chemical composition of 12035.

reference	Gast70	Hubbard71	Weismann75	Cuttitta71	Compston71	Tatsumoto71
weight		141 mg	141 mg			
SiO ₂ %				43.17	(c)	
TiO ₂				2.28	(c)	
Al ₂ O ₃				8.03	(c)	
FeO				22.2	(c)	
MnO			0.27	(b) 0.29	(c)	
MgO				15.49	(c)	
CaO				8.08	(c)	
Na ₂ O				0.21	(c)	
K ₂ O	0.044	(a) 0.044	(a) 0.044	(a)	0.054	(c)
P ₂ O ₅				0.06	(c)	
S %				0.05	(c)	
<i>sum</i>						
Sc ppm			42	(b)		
V			204	(b) 130	(c)	
Cr			4540	(b) 3360	(c)	
Co			71	(b) 52	(c)	
Ni			101	(b) 33	(c)	
Cu			6.5	(b) 2	(c)	
Zn						
Ga			4.2	(b) 1.8	(c)	
Ge ppb						
As						
Se						
Rb	0.682	(a) 0.689	(a) 0.689	(a) 1.2	(b) 0.83	(c)
Sr		84.3	(a) 84.3	(a) 65	(b) 100.4	(c)
Y				36	(b) 29	(c)
Zr				88	(b) 81	(c)
Nb					4	(c)
Mo						
Ru						
Rh						
Pd ppb						
Ag ppb						
Cd ppb						
In ppb						
Sn ppb						
Sb ppb						
Te ppb						
Cs ppm						
Ba	47.2	(a) 47.2	(a) 47.2	(a) 51	(b) 45	(c)
La		3.87	(a) 3.87	(a)	3	(c)
Ce	11.5	(a) 11.5	(a) 11.5	(a)	6	(c)
Pr						
Nd	8.91	(a) 8.91	(a) 8.91	(a)		
Sm	3.22	(a) 3.22	(a) 3.22	(a)		
Eu	0.751	(a) 0.751	(a) 0.751	(a)		
Gd	4.32	(a) 4.3	(a) 4.3	(a)		
Tb						
Dy	5.07	(a) 5.07	(a) 5.09	(a)		
Ho						
Er	3.09	(a) 3.09	(a) 3.09	(a)		
Tm						
Yb	3.04	(a) 3.04	(a) 2.71	(a) 4.7	(b)	
Lu	0.423	(a) 0.423	(a) 0.423	(a)		
Hf			0.15	(a)		
Ta						
W ppb						
Re ppb						
Os ppb						
Ir ppb						
Pt ppb						
Au ppb						
Th ppm				0.801	0.682	(a)
U ppm				0.24	0.199	(a)

technique (a) IDMS, (b) mixed microchem. XRF and emission spec., (c) XRF



List of Photo #'s for 12035

S69-61249-61256	group
S69-63158-63165	color
S70-18927-18934	B & W, mug ,9
S70-44328	best color
S70-44329	
S70-44174-44179	color
S70-45626	TS reflected
S70-45637	TS transmitted
S70-49427-49430	TS color
S70-49465-49466	TS color

Compston W., Berry H., Vernon M.J., Chappell B.W. and Kay M.J. (1971) Rubidium-strontium chronology and chemistry of lunar material from the Ocean of Storms. *Proc. 2nd Lunar Sci. Conf.* 1471-1485.

Cuttitta F., Rose H.J., Annell C.S., Carron M.K., Christian R.P., Dwornik E.J., Greenland L.P., Helz A.P. and Ligon D.T. (1971) Elemental composition of some Apollo 12 lunar rocks and soils. *Proc. 2nd Lunar Sci. Conf.* 1217-1229.

Gast P.W. and Hubbard N.J. (1970c) Rare earth abundances in soil and rocks from the Ocean of Storms. *Earth Planet. Sci. Lett.* **10**, 94-101.

Hubbard N.J. and Gast P.W. (1971) Chemical composition and origin of nonmare lunar basalts. *Proc. 2nd Lunar Sci. Conf.* 999-1020.

James O.B. and Wright T.L. (1972) Apollo 11 and 12 mare basalts and gabbros: Classification, compositional variations and possible petrogenetic relations. *Geol. Soc. Am. Bull.* **83**, 2357-2382.

Neal C.R., Hacker M.D., Snyder G.A., Taylor L.A., Liu Y.-G. and Schmitt R.A. (1994a) Basalt generation at the Apollo 12 site, Part 1: New data, classification and re-evaluation. *Meteoritics* **29**, 334-348.


References for 12035

Brett R., Butler P., Meyer C., Reid A.M., Takeda H. and Williams R. (1971) Apollo 12 igneous rocks 12004, 12008, 12009 and 12022: A mineralogical and petrological study. *Proc. 2nd Lunar Sci. Conf.* 301-317.

Butler P. (1972b) Compositional characteristics of olivines from Apollo 12 samples. *Geochim. Cosmochim. Acta* **36**, 773-785.

Burnett D.S., Monnin M., Seitz M., Walker R. and Yuhas D. (1971) Lunar astrology – U-Th distributions and fission-track dating of lunar samples. *Proc. 2nd Lunar Sci. Conf.* 1503-1519.

THE CHIPPING OF LUNAR ROCK 12035,9



Neal C.R., Hacker M.D., Snyder G.A., Taylor L.A., Liu Y.-G. and Schmitt R.A. (1994b) Basalt generation at the Apollo 12 site, Part 2: Source heterogeneity, multiple melts and crustal contamination. *Meteoritics* **29**, 349-361.

Papanastassiou D.A. and Wasserburg G.J. (1971a) Lunar chronology and evolution from Rb-Sr studies of Apollo 11 and 12 samples. *Earth Planet. Sci. Lett.* **11**, 37-62.

Papike J.J., Hodges F.N., Bence A.E., Cameron M. and Rhodes J.M. (1976) Mare basalts: Crystal chemistry, mineralogy and petrology. *Rev. Geophys. Space Phys.* **14**, 475-540.

Reid J.B. (1971) Apollo 12 spinels as petrogenetic indicators. *Earth Planet. Sci. Lett.* **10**, 351-356.

Reid A.M., Meyer C., Harmon R.S. and Brett R. (1970c) Metal grains in Apollo 12 igneous rocks. *Earth Planet. Sci. Lett.* **9**, 1-5.

Roedder E. and Weiblen P.W. (1971) Petrology of silicate melt inclusions, Apollo 11 and Apollo 12 and terrestrial equivalents. *Proc. Second Lunar Sci. Conf.* 507-528.

Schnetzler C.C. and Philpotts J.A. (1971) Alkali, alkaline earth, and rare earth element concentrations in some Apollo 12 soils, rocks, and separated phases. *Proc. 2nd Lunar Sci. Conf.* 1101-1122.

Tatsumoto M., Knight R.J. and Doe B.R. (1971) U-Th-Pb systematic of Apollo lunar samples. *Proc. 2nd Lunar Sci. Conf.* 1521-1546.

Walker D., Kirkpatrick R.J., Longhi J. and Hays J.F. (1976) Crystallization history of lunar picritic basalt sample 12002: Phase-equilibria and cooling-rate studies. *Geol. Soc. Am. Bull.* **87**, 646-656.

Warner J. (1970) Apollo 12 Lunar Sample Information. NASA TR R-353. JSC (catalog)

Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample Data Generated by the Gast, Nyquist and Hubbard Lunar Sample PI-Ships. Unpublished. JSC