

**78221**  
Soil (under boulder)  
345 grams



Figure 1: Lunar soil sample 78220 was collected from beneath the boulder (78235). AS17-146-22371.



Figure 3: Surface photo after boulder (figure 1) rolled away and soil sample 78220 collected from beneath boulder. AS17-142-21705.




Figure 2: Map of station 8, Apollo 17.




Figure 4: Composition of lunar soils showing 78221.

**Introduction**

Lunar soil sample 78221 was chosen a “reference soil” for highland initiative (Papike et al. 1982; Simon et al. 1981). It was collected at station 8, Apollo 17, on the bottom slope of the Sculptured Hills (Wolfe et al. 1981). It is a very mature soil from beneath the norite boulder (78235-78255) after it was rolled away. It was found to contain a significant amount of basalt.

**Petrography**

The maturity of 78221 is  $I_s/FeO = 93$  and the average grain size is 43 microns (Morris 1978, Graf 1993). This is a very mature soil with agglutinate content 57 % (Heiken and McKay 1974).

Both Heiken and McKay (1974) and Simon et al. (1981) determined the mineralogical mode (see table), but they studied different grain sizes.




Figure 5: Carbon content and maturity index for 78221 and other lunar soils.

Meyer (1973) cataloged the coarse-fine particles and Blanchard et al. (1975) studied the 1 – 2 mm fraction. Blanchard et al. reported on 31 “mare basalt”, 22 “glassy breccias”, and 12 “highland rocks”, but gave few details.

### Mineralogy

Simon et al. (1981) determined the composition of olivine, pyroxene and plagioclase from two different size fractions of 78221, finding a significant contribution from mare basalt, but also some Mg-rich grains (figure 8).

### Chemistry

The chemical composition was determined by Duncan et al. (1974), Blanchard et al. (1975), Laul et al. (1981) and Korotev and Kremser (1992). It is similar to the other station 8 soils and is depleted in the heavy rare earth elements (Gd – Lu) compared with mare and highland soils (figure 6). Laul et al. (1981) also determined the chemical composition of different grain





Figure 6: Normalized rare-earth-element diagram for 78221 compared with that of mare and highland soils.

size separates (figure 9). They found increased amounts of trace elements in the finest fraction.

Moore et al. (1974) determined 190 ppm carbon (figure 5).



average grain size = 43 microns




Figure 7: Grain size distribution for 78220 (Graf 1993, data from King)

### Modal content of soil 78221

From	Heiken and McKay 1974	Simon et al. 1981
	90-150 microns	90-1000 microns
Agglutinates	57%	46.6
Basalt	1	5.7
Breccia	13.3	12
Anorthosite	1.3	2.2
Norite		
Gabbro	0.3	
Plagioclase	5	9.9
Pyroxene	12.5	9.8
Olivine	1.7	
Ilmenite	1	0.4
Glass other	7	6.5




Figure 8: Pyroxene and olivine compositions for two different grain size fraction of 78220 (Simon et al. 1981).

**Cosmogenic isotopes and exposure ages**

This soil was shielded by the meter-sized boulder that sat on top of it. However, since the soil is mature it would have had a previous radiation history.





Figure 9: Normalized rare-earth-element diagram for 78220 and its various grain size fractions (Laul et al. 1981).



**Table 1. Chemical composition of 78221.**

reference	Korotev92		Duncan74	Laul81	<20 micron	78222 Blanchard75 1 - 2 mm ave. 39			
<i>weight</i>									
SiO2 %			43.67	(b) 43	43.9	(a)			
TiO2			3.84	(b) 4.2	3.9	(a)			
Al2O3			17.13	(b) 17	19	(a)			
FeO	12	11.9	(a) 11.68	(b) 12.6	11.9	(a) 11.9	12	(a)	
MnO			0.157	(b) 0.163	0.135	(a) 0.17	0.19	(a)	
MgO			10.55	(b) 11	9.3	(a)			
CaO			11.79	(b) 12.4	12.4	(a)			
Na2O	0.376	0.378	(a) 0.37	(b) 0.37	0.4	(a) 0.4	0.49	(a)	
K2O			0.092	(b) 0.1	0.12	(a)			
P2O5			0.08	(b)					
S %			0.088	(b)					
<i>sum</i>									
Sc ppm	34.6	34.3	(a)		36.3	29.3	(a) 32	34.4	(a)
V			68	(b) 70	65	(a)			
Cr	2370	2270	(a) 2196	(b) 2395	2190	(a) 2220	2550	(a)	
Co	35.9	37	(a) 34	(b) 40	36.5	(a) 37.8	31	(a)	
Ni	280	230	(a) 221	(b) 200	600	(a) 260	280	(a)	
Cu			5.8	(b)					
Zn			25.6	(b)					
Ga									
Ge ppb									
As									
Se									
Rb			2.6	(b)					
Sr	140	170	(a) 147	(b) 160	170	(a)			
Y			45	(b)					
Zr	180	180	(a) 173	(b)					
Nb			13.3	(b)					
Mo									
Ru									
Rh									
Pd ppb									
Ag ppb									
Cd ppb									
In ppb									
Sn ppb									
Sb ppb									
Te ppb									
Cs ppm									
Ba	118	106	(a) 109	(b) 120	170	(a)			
La	8.64	9.01	(a)	8.1	11.4	(a) 8.5	11.1	(a)	
Ce	23.2	25.4	(a)	24	30	(a) 23.9	27	(a)	
Pr									
Nd		21	(a)	16	22	(a)			
Sm	5.83	6.06	(a)	5.46	6.9	(a) 6	5.64	(a)	
Eu	1.31	1.34	(a)	1.25	1.5	(a) 1.23	1.36	(a)	
Gd									
Tb	1.35	1.4	(a)	1.3	1.5	(a) 1.4	1.43	(a)	
Dy				8.4	10	(a)			
Ho									
Er									
Tm				0.72	0.81	(a)			
Yb	4.74	5.11	(a)	4.7	5.4	(a) 5.34	4.86	(a)	
Lu	0.68	0.693	(a)	0.7	0.79	(a) 0.73	0.7	(a)	
Hf	4.89	4.97	(a)	4.7	5	(a) 4.7	4.6	(a)	
Ta	0.76	0.76	(a)	0.86	0.93	(a) 1	0.9	(a)	
W ppb									
Re ppb									
Os ppb									
Ir ppb	8.2	11.5	(a)						
Pt ppb									
Au ppb	5	3	(a)						
Th ppm	1.1	1.3	(a)	1.6	1.85	(a)			
U ppm	0.22	0.34	(a)	0.4	0.4	(a)			

*technique: (a) INAA, (b) XRF*

## References for 78221

- Blanchard D.P., Krotev R.L., Brannon J.C., Jacobs J.W., Haskin L.A. Reid A.M., Donaldson C. and Brown R.W. (1975) A geochemical and petrographic study of 1-2 mm fines from Apollo 17. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 2321-2342.
- Butler P. (1973) Lunar Sample Information Catalog Apollo 17. Lunar Receiving Laboratory. MSC 03211 Curator's Catalog. pp. 447.
- Butler J.C. and King E.A. (1974) Analysis of the grain size-frequency distributions of lunar fines. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 829-841.
- Cripe J.D. and Moore C.B. (1976) Extralunar sulfur in Apollo 16 and 17 lunar fines. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 469-479.
- Duncan A.R., Erlank A.J., Willis J.P., Sher M.K. and Ahrens L.H. (1974) Trace element evidence for a two-stage origin of some Titaniferous mare basalts. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1147-1158.
- Gibson E.K. and Moore G.W. (1974) Sulfur abundances and distributions in the valley of Taurus-Littrow. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1823-1838.
- Green G.M., King D.T., Banholzer G.S. and King E.A. (1975) Size and model analyses of fines and ultrafines from some Apollo 17 samples. *Proc. 6<sup>th</sup> Lunar Sci. Conf.* 517-528.
- Graf J.C. (1993) Lunar Soils Grain Size Catalog. NASA Reference Pub. 1265, March 1993
- Heiken G.H. (1974) A catalog of lunar soils. JSC Curator
- Heiken G.H. (1975) Petrology of lunar soils. *Rev. Geophys. Space Phys.* **13**, 567-587.
- Heiken G.H. and McKay D.S. (1974) Petrology of Apollo 17 soils. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 843-860.
- Korotev R.L. and Kremser D. (1992) Compositional variations in Apollo 17 soils and their relationships to the geology of the Taurus-Littrow site. *Proc. 22<sup>nd</sup> Lunar Planet. Sci. Conf.* 275-301.
- Laul J.C., Papike J.J. and Simon S.B. (1981) The lunar regolith: Comparative studies of the Apollo and Luna 16, 20 and 24. *Proc. 12<sup>th</sup> Lunar Planet. Sci. Conf.* 389-407.
- LSPET (1973a) Apollo 17 lunar samples : Chemical and petrographic description. *Science* **182**, 659-690.
- LSPET (1973c) Preliminary examination of lunar samples. Apollo 17 Preliminary Science Report. NASA SP-330, 7-1—7-46.
- McKay D.S., Fruland R.M. and Heiken G.H. (1974) Grain size and the evolution of lunar soils. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 887-906.
- Meyer C. (1973) Apollo 17 Coarse Fines (4-10 mm) Sample Location, Classification and Photo Index. Curator Report. pp. 182.
- Moore C.B., Lewis C.F. and Cripe J.D. (1974a) Total carbon and sulfur contents of Apollo 17 lunar samples. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1897-1906.
- Morris R.V. (1976) Surface exposure indices of lunar soils: A comparative FMR study. *Proc. 7<sup>th</sup> Lunar Sci. Conf.* 315-335.
- Morris R.V., Score R., Dardano C. and Heiken G. (1983) Handbook of Lunar Soils. Two Parts. JSC 19069. Curator's Office, Houston
- Morris R.V. (1978) The surface exposure (maturity) of lunar soils: Some concepts and Is/FeO compilation. *Proc. 9<sup>th</sup> Lunar Sci. Conf.* 2287-2297.
- Morris R.V. (1980) Origins and size distribution of metallic iron particles in the lunar regolith. *Proc. 11<sup>th</sup> Lunar Planet. Sci. Conf.* 1697-1712.
- Papike J.J., Simon S.B. and Laul J.C. (1982) The lunar regolith: Chemistry, Mineralogy and Petrology. *Rev. Geophys. Space Phys.* **20**, 761-826.
- Schonfeld E. (1974) The contamination of Lunar highlands rocks by KREEP: Interpretation by mixing models. *Proc. 5<sup>th</sup> Lunar Sci. Conf.* 1269-1286.
- Simon S.B., Papike J.J. and Laul J.C. (1981) The lunar regolith: Comparative studies of the Apollo and Luna sites. *Proc. 12<sup>th</sup> Lunar Planet. Sci. Conf.* 371-388.
- Wolfe E.W., Bailey N.G., Lucchitta B.K., Muehlberger W.R., Scott D.H., Sutton R.L and Wilshire H.G. (1981) The geologic investigation of the Taurus-Littrow Valley: Apollo 17 Landing Site. US Geol. Survey Prof. Paper, 1080, pp. 280.