

QUE94201 – 12.02 grams Depleted Basaltic Shergottite



Figure 1. Photograph illustrating broken interior surface of Martian meteorite QUE94201. Scale is in mm. Cube is 1 cm. (NASA # S96-00376)

Introduction

QUE94201 was found on a glacial moraine where it “looked different”. A search for other pieces proved unsuccessful (Lofgren, personal communication).

Five sides of QUE94201 are rounded with remnant fusion crust, while one side appears freshly broken (figure 1). The interior is coarse-grained, crystalline and glassy (Score and Mason 1995). “Mafic-rich areas” (probably shock-melted glass), as large as 5 x 4 mm, were noted during preliminary examination. In thin section, the sample is made up of subequal amounts of homogeneous maskelynite laths and variable interstitial pyroxene. Maskelynite laths are up to 3.6 mm long.

QUE94201 is a basalt apparently similar to the dark, mottled lithology (DML) of Zagami (McSween *et al.* 1996) as well as “lithology B” of EETA 79001 (Mikouchi *et al.* 1998). However, the phosphorous content of QUE94201 is high and the REE pattern is strongly depleted in light rare earth elements. The extreme

zoning in pyroxene in QUE94201 indicates that it cooled quickly from magmatic temperatures. The Fe-Ti oxide compositions indicate that this basalt formed under more reducing conditions than the other shergottites (McSween *et al.* 1996; Herd *et al.* 2001d; Kring *et al.* 2003). Karner *et al.* (2007, 2008) have used Cr and V partitioning between pyroxene and melt, and the valance state of V to exactly determine the reducing conditions.

Kring *et al.* (2003) relate QUE94201 to the lherzolithic shergottites, because of the LREE depletion, but note that it was a basaltic liquid before it crystallized. Warren *et al.* (1999) find that QUE94201 analyses are very low in Ni and Ir content, indicating that it was formed from a magma-source in the interior of Mars, and unrelated to meteorite bombardment.

The age of QUE94201 is 327 m.y. with an exposure to cosmic ray for 2.2 m.y.




Figure 2. Photomicrograph of thin section of QUE94201,4 illustrating coarse basaltic texture (intergrown plagioclase and pyroxene). Field of view is 2.2 mm.

Petrography

Harvey *et al.* (1996) described QUE94201 as a “coarse-grained basalt, consisting of subhedral Fe-rich pigeonite and maskelynite”. Kring *et al.* (2003) not that it was a “bulk melt”, meaning that its composition can be used to infer the magmatic source region. Most of the pyroxene and maskelynite grains exceed 1 mm in length (up to 3 mm) and are somewhat elongated (figures 2; 7). QUE94201 contains relatively high proportions of maskelynite and phosphate when compared with the other shergottites. The texture is nicely illustrated in figure 7. Melt inclusions are found in the relict plagioclase (Kring *et al.* 2003), but no melt inclusions were found in the pyroxene.

The pyroxenes in QUE94201 are complexly zoned (McKay *et al.* 1996; Mikouchi *et al.* 1996; 1998 and McSween and Eisenhour 1996). The Mg-rich pigeonite cores are mantled by Mg-rich augite, which is, in turn, rimmed by Fe-rich pigeonite and strongly zoned to pyroxferroite (with some hedenbergite at the edge). None of the cores appear to be cumulate phases, as was the case for Shergotty, Zagami and EETA79001B. Some of the pyroxenes in QUE94201 are also found to be sector zoned.

Interstitial to the pyroxene and shocked plagioclase, are a number of late-stage phases including large Fe-Ti oxides (ulvöspinel, rutile, ilmenite), whitlockite and large “pockets” of mesostasis similar to the “DN pockets” of Zagami (McCoy *et al.* 1995). These “pockets” contain an intergrowth of silica and fayalite, as well as, maskelynite, whitlockite, Fe-Ti oxides, sulfides, minor augite, chlorapatite and a Zr-rich phase, probably baddelyite (McSween *et al.* 1996). Fayalite-silica intergrowths are also found in the cores of large skeletal

Mineralogical Mode for QUE94201

	Harvey <i>et al.</i> (1998)	Mikouchi <i>et al.</i> (1996)
Pyroxene	44 vol. %	43
Maskelynite	46	42
Opaque	2	4
Phosphate	4	6
Mesostasis	4	5




Figure 3. Composition diagram for pyroxene and olivine in QUE94201. Data are from Kring *et al.* (1996), McKay *et al.* (1996), McSween and Eisenhour (1996) and Mikouchi *et al.* (1996). Olivine is pure fayalite.

phosphate grains adjacent to these pockets (Harvey *et al.* 1996). Aramovich *et al.* (2002) have carefully studied the symplectite intergrowths associated with merrillite grains in QUE94201 and conclude that the early formation of Ca and Mg-rich merrillite lead to the formation of metastable pyroxferroite and ferrosilite, which then broke down to become the fine-grained intergrowths.

Shock features include maskelynite, mosaicism in pyroxene and large pockets of glass formed *in-situ*. The shock-melted glass is rich in phosphorous (Mikouchi *et al.* 1998). Papike *et al.* (2009) compare QUE94201 with the other shergottites.

Mineral Chemistry

Pyroxene: Pyroxene zoning is extreme (figures 3, 8), including sector zoning in the cores (Kring *et al.* 1996; Mikouchi *et al.* 1996, 1998; McKay *et al.* 1996; McSween and Eisenhour 1996). Papike (2005 onward) has been obsessed with these pyroxenes. Harvey *et al.* (1996) report pigeonite zoning to Fs_{85} . Wadhwa and Crozaz (1996), McSween *et al.* (1996) and Wadhwa *et al.* (1998) have determined the REE patterns of the pyroxenes and suggested that merrillites began crystallizing relatively early. The Eu anomaly indicates reducing conditions.

“Pyroxferroite”: An analysis of “pyroxferroite” is given in Mikouchi *et al.* (1998), however, Aramovich *et al.* (2002) point out that the reported composition ($Wo_{37}En_3Fs_{60}$) is too Ca-rich for it to be termed “pyroxferroite”.




Figure 4. Normalized rare earth element diagram for QUE94201 compared with that of Shergotty.

Maskelynite: Plagioclase (An_{66-52}) crystallized throughout the crystallization sequence (McSween and Eisenhour 1996; Mikouchi *et al.* 1999). The cores are the most An-rich among the basaltic Martian meteorites. It has been shocked to maskelynite. Plagioclase is found to include melt inclusions (Kring *et al.* 2003).

Phosphates: QUE94201 contains more phosphates than other SNC meteorites. Whitlockite (merrillite) has been studied by Wadhwa and Crozaz (1996) and found to have a more extreme depletion of LREE than for any other shergottite. Mikouchi *et al.* (1998) analyzed merrillite up to 3 mm long. Mikouchi *et al.* (1996, 1998), McSween *et al.* (1996) and Leshin *et al.* (1996) also report minor chlorapatite in the mesostasis. Aramovich *et al.* (2002) show that the formation of merrillite is often surrounded by symplectite intergrowths.

Silica: Silica occurs as distinctive intergrowth with fayalite in “patches” up to 1 mm in-between pyroxene and plagioclase grains (Harvey *et al.* 1996). Silica was also reported as an alteration product by Wentworth and Gooding (1991).

Olivine: Fayalite (Fa_{96-99}) occurs as a fine dendritic intergrowth with silica (Harvey *et al.* 1996).

Opaques: Large grains of ilmenite are the major opaque phase. Analyses for ilmenite and ulvöspinel are reported in Kring *et al.* (1996), McSween *et al.* (1996), Mikouchi *et al.* (1998) and Herd *et al.* (2001). Chromite has been found in Mg-rich pigeonite core (Mikouchi *et al.*). Ulvöspinel exhibits subsolidus

Table 1a. Chemical composition of QUE94201.

	Warren96	Dreibus96	Kring 96	Kring 96	Mikouchi96	Mittlefehldt 96	Warren 97
weight	305 mg	179.8 mg	250mg	fusion crust	fusion crust	52.74 mg	305 mg
SiO ₂ %	47.06 (a)			43.5 (d)	44.3 (d)		48.00 (d)
TiO ₂	1.95 (a)	1.8 (b)	1.7 (b)	1.81 (d)	2 (d)		1.98 (b)
Al ₂ O ₃	9.64 (a)	12 (b)	11.1 (b)	7.46 (d)	7 (d)		9.82 (b)
FeO	18.65 (a)	18.3 (b)	18.3 (b)	24.2 (d)	21 (d)	20.0 (b)	19.16
MnO	0.48 (a)	0.436 (b)	0.44 (b)	0.63 (d)	0.6 (d)		0.47 (b)
CaO	11.3 (a)	11.3 (b)		10.9 (d)	11.2 (d)	10.7 (b)	11.48
MgO	6.3 (a)	6.2 (b)		6.04 (d)	6.4 (d)		6.3 (d)
Na ₂ O	1.39 (a)	1.75 (b)		1.16 (d)	1.1 (d)	1.53 (b)	1.39
K ₂ O	0.038 (a)	0.052 (b)		0.04 (d)			0.04
P ₂ O ₃				2.77 (d)	3.4 (d)		
sum	96.81			98.51	97		98.64
Li ppm							
C							
F		40 (b)					
S							
Cl		91 (b)					
Sc	49 (b)	46.6 (b)				51.5 (b)	49.0 (b)
V	124 (b)	103 (b)					124
Cr	1030 (b)	890 (b)					1010
Co	24.4 (b)	22.8 (b)				25.9 (b)	24.4
Ni	<40 (c)	<20 (b)					<40
Cu							
Zn	108 (c)					130	108
Ga	26 (b)	27.1 (b)					25.9
Ge							
As		0.77 (b)					
Se							
Br		0.35 (b)		Borg 97	Borg 97	0.38	
Rb		(b)		0.518 (f)	0.691 (f)		<6
Sr	59 (b)	80 (b)		41.3 (f)	49.8 (f)	70	59
Y		31.2 (e)					
Zr	94 (b)	97.1 (e)				80	94
Nb		0.68 (e)					
Mo							
Pd ppb							
Ag ppb							
Cd ppb							
In ppb							
Sb ppb							
Te ppb							
I ppm		4.6 (b)					
Cs ppm							<0.12
Ba	<41 (b)	<15 (b)					<41
La	0.44 (b)	0.35 (b)				0.31	0.44
Ce	1.63 (b)	1.3 (b)				1.0	1.63
Pr				Borg 97			
Nd	2.4 (b)	1.9 (b)		1.482 (f)			2.36
Sm	2.55 (b)	2.02 (b)		1.233 (f)		1.92	2.55
Eu	1.09 (b)	0.99 (b)				0.9	1.09
Gd		4.3 (b)					
Tb	0.93 (b)	0.802 (b)				0.78	0.93
Dy	6.1 (b)	5.53 (b)					
Ho		1.19 (b)					
Er							
Tm							
Yb	3.5 (b)	3.09 (b)				3.02	
Lu	0.54 (b)	0.455 (b)				0.42	0.54
Hf	3.4 (b)	3.42 (b)				4.2	3.4
Ta	<0.08 (b)	0.023 (b)				0.03	<0.08
W ppb							
Re ppb							
Os ppb							
Ir ppb	<2.4 (b)	<3 (b)					<2.4
Au ppb		<1.5 (b)					<0.5
Tl ppb							
Bi ppb							
Th ppm	<0.09 (b)	0.05 (e)					<0.09
U ppm		0.0125 (e)					<0.2

technique: (a) emp fused bead, (b) INAA, (c) RNAA, (d) emp, (e) spark source mass spec., (f) isotope dilution mass spec.

Table 1b. Chemical composition of QUE94201.

reference	Lodders 98	Wang 98	Warren 99	Blichert-Toft 99	Kring 2003						
weight	average		285 mg.	50 mg.	48 mg.	57 mg.	47.4 mg	81.32	125.87	254.6	
SiO ₂	47.9		47.92								
TiO ₂	1.84		1.98	(a)			1.6	1.82	1.67	1.7 (a)	
Al ₂ O ₃	11		9.82	(a)			15.17	11.7	10.94	11.98 (a)	
FeO	18.5		19.17	(a)			15.37	18.65	18.01	17.73 (a)	
MnO	0.45		0.47	(a)			0.38	0.44	0.45	0.44 (a)	
CaO	11.4		11.47	(a)			10.77	10.35	10.65	10.63 (a)	
MgO	6.25		6.3	(a)			5.47	4.81	6.63	5.8 (a)	
Na ₂ O	1.58		1.37	(a)			2.13	1.86	1.79	1.87 (a)	
K ₂ O	0.045		0.04	(a)			0.052	0.061	0.077	0.067 (a)	
P ₂ O ₅											
sum											
Li ppm											
Sc	48		49	(a)			38.5	45.8	48.8	45.9 (a)	
V	113		124	(a)			84	92	104	96 (a)	
Cr	950		1010	(a)			792	809	1054	927 (a)	
Co	24	13.7	(b) 24.4	(a)			20.22	23.1	24.2	23.1 (a)	
Ni	<20		6.6	(b)			150	120	<80	<106 (a)	
Cu											
Zn	110	87.4	(b) 90	(b)			65	82	84	80 (a)	
Ga	27	20.1	(b) 25.9	(a)			27.6	30.5	25	27.2 (a)	
Ge			1.95	(b)							
As	0.77		<0.6								
Se		0.473	(b)				2.9	2.8	1.8	2.3 (a)	
Br	0.35		0.5	(a)							
Rb		0.545	(b)				<13		8.4	8.4 (a)	
Sr	70		59	(a)							
Y	31										
Zr	100		94	(a)			118	159	107	126 (a)	
Nb	0.68										
Mo											
Pd ppb											
Ag ppb		3.46	(b)								
Cd ppb		52.6	(b) <50	(b)							
In ppb		76.7	(b)								
Sb ppb		5.3	(b)								
Te ppb		4.1	(b)								
I ppm	4.6										
Cs ppm		0.032	(b) <0.120	(a)							
Ba	<15										
La	0.4		0.44	(a)			0.35	0.45	0.46	0.44 (a)	
Ce	1.47		1.63	(a)			<1.8	1.62	1.19	1.36 (a)	
Pr											
Nd	2.2		2.36	(a)			<5	3.8	3.1	3.4 (a)	
Sm	2.3		2.55	(a)			1.747	2.53	2.436	2.34 (a)	
Eu	1.04		1.09	(a)			1.086	1.208	1.144	1.154 (a)	
Gd	4.3										
Tb	0.87		0.93	(a)			0.76	1.034	1.026	0.979 (a)	
Dy	5.8		6.1	(a)							
Ho	1.19		1.32	(a)			0.95	1.35	1.23	1.22 (a)	
Er											
Tm							0.45	0.58	0.51	0.52 (a)	
Yb	3.3		3.5	(a)			2.7	3.77	3.5	3.44 (a)	
Lu	0.5		0.54	(a)	0.403	0.389	0.727	(c) 0.383	0.556	0.496	0.494 (a)
Hf	3.41		3.4	(a)	2.99	3.17	4.28	(c) 2.76	3.9	2.61	3.05 (a)
Ta	0.023		<0.08	(a)							
W ppb							<1.1	<0.8	0.54	0.54 (a)	
Re ppb			0.0028	(b)			60	42	29	39 (a)	
Os ppb			0.0051	(b)							
Ir ppb	<3		0.012	(b)			<12	6	5	5 (a)	
Au ppb	<0.5	0.145	(b) 0.041	(b)			<7	5	<4	(a)	
Tl ppb		20.9	(b)								
Bi ppb		1.54	(b)								
Th ppm	<0.05		<0.09	(a)							
U ppm	<0.05	0.023	(b) <0.2	(a)							
technique	(a) INAA, (b) RNAA, (c) IDMS										




Figure 5. Rb-Sr isochron for QUE94201 as reported by Borg *et al.* (1997).




Figure 6. Sm-Nd isochron for QUE94201 as determined by Borg *et al.* (1997).

“oxyexsolution” of ilmenite lamellae on the order of 300-500 nm wide (Herd *et al.*)

Sulfide: The sulfide phase is pyrrhotite (McKay *et al.* 1996; McSween and Eisenhour 1996).

Glass: QUE94201 contains abundant pockets of shock-melted glass. This melt contains up to 7 % P, probably due to preferential melting of the abundant phosphates (Mikouchi *et al.* 1996, 1998). K-poor and K-rich feldspar glasses have also been reported in the mesostasis by Mikouchi *et al.*

Salts: Fe-K-sulfates are sometimes observed rimming Fe-sulfides (Harvey *et al.* 1996). The salts have been studied in detail by Wentworth and Gooding (1996). They found that “carbonates are conspicuously absent.”

Baddelyite: ZrO₂ is reported in the mesostasis (McSween *et al.* 1996).

Whole-rock Composition

The chemical composition of QUE94201 has been determined and reported by Dreibus *et al.* (1996), Warren and Kallemeyn (1996, 1997), Mittlefehldt and Lindstrom (1996), Warren *et al.* (1999) and Kring *et al.* (2003). The sample has very high phosphorous content (table 1). This is also reflected in the analysis of the fusion crust and modal mineralogy of the thin sections (Kring *et al.* 1996; Mikouchi *et al.* 1996). Note that Ni and Ir are rather low (6 and 0.012 ppm, Warren *et al.*)(but apparently high in Kring *et al.* 2003).

QUE94201 is a basalt that is greatly depleted in LREE

(figure 4). Kring *et al.* (2003) make a careful effort to compute the major element composition. Key to this computation is knowledge of P and Si contents as determined from fused bead analyses (Mittlefehldt and Lindstrom 1996) and fusion crust (Kring *et al.* 1996).

Radiogenic Isotopes

Borg *et al.* (1996, 1997) reported a Rb-Sr age ($\lambda_{Rb} = 1.402 \times 10^{-11} \text{ year}^{-1}$) of $327 \pm 12 \text{ m.y.}$ with initial $^{87}\text{Sr}/^{86}\text{Sr}$ ratio of 0.701298 ± 14 (figure 5). This low I_{Sr} ratio indicates that the source region (Martian mantle) was depleted in Rb. The Sm-Nd age of $327 \pm 19 \text{ m.y.}$ with $\epsilon_{Nd} = 47.6 \pm 1.7$ is concordant with the Rb-Sr age (figure 6). Dreibus *et al.* (1996b) reported a K/Ar age of 1.33 b.y. and Bogard and Garrison (1999) reported 730 m.y. by Ar/Ar plateau.

Cosmogenic Isotopes and Exposure Ages

From cosmic-ray produced ^3He , ^{21}Ne and ^{38}Ar , Eugster *et al.* (1996) computed an exposure age for QUE94201 of $2.4 \pm 0.6 \text{ m.y.}$ and concluded that QUE94201 was “ejected from Mars simultaneously with the other basaltic shergottites - Shergotty and Zagami”. Nishiizumi and Caffee (1996) found the ^{10}Be concentrations gave a cosmic-ray exposure age of $2.6 \pm 0.5 \text{ m.y.}$ for an assumed 4π irradiation geometry. Garrison and Bogard (1998) determined a cosmic ray exposure age of $2.7 \pm 0.6 \text{ m.y.}$ Terribillini *et al.* (2000) and Eugster *et al.* (2002) used ^{81}Kr to determine exposure ages of 2.10 ± 0.25 and $2.22 \pm 0.35 \text{ m.y.}$ (respectively). Dreibus *et al.* (1996b) and Swindle *et al.* (1996) have also reported exposure ages.

Nishiizumi and Caffee (1996) found that the terrestrial age ($0.29 \pm 0.05 \text{ m.y.}$ obtained from ^{36}Cl) of QUE94201

is longer than for other Antarctic shergottites. Jull *et al.* (1997) found that ^{14}C activity was nil, consistent with an old terrestrial age.

Schnabel *et al.* (2001) reported ^{10}Be , ^{26}Al and ^{53}Mn activities of 11.9 ± 0.2 , 63.4 ± 6.9 and 162 ± 11 dpm/kg respectively. Kring *et al.* (2003) sum the terrestrial age (0.29 m.y.), with the cosmic ray exposure age (2.6 m.y.) to get an ejection age from Mars of 2.9 m.y.

Other Isotopes

Oxygen isotopes were reported by Clayton and Mayeda (1996) and Franchi *et al.* (1997)(figure I-3).

Leshin *et al.* (1996, 2000) determined that the hydrogen in six “apatite” grains in QUE94201 has a high D/H ratio ($\delta\text{D} = 1700$ to 3500 ‰), probably from the Martian hydrosphere. Boctor *et al.* (2001) also determined the isotopic composition of hydrogen in feldspathic glass.

Grady *et al.* (1996) reported that the carbon released from 450 to 600°C was isotopically light ($\delta^{13}\text{C} \sim -24.2$ ‰).

Eugster *et al.* (1996), Swindle *et al.* (1996) and Garrison and Bogard (1998) determined the contents and isotopic ratios of rare gases (Ne, Ar, Kr, Xe) in QUE94201 and found them typical of other shergottites. Small amounts of ^{21}Ne produced by energetic solar protons may be present in QUE94201 (Garrison and Bogard 1998). Garrison and Bogard’s (1998) study was for unmelted mineral phases.

Blichert-Toft *et al.* (1998, 1999) found a very large Hf isotopic anomaly - to match the large Nd and W isotopic anomalies as determined by Borg *et al.* (1997) and Lee and Halliday (1997). Gafney *et al.* (2006, 2007) studied the U-Pb systematics and determined a low U/Pb ratio for the Martian mantle. These isotopic anomalies have been preserved in the source region of Martian basalts, since the initial early formation of the crust of Mars!

Experimental Studies

Since QUE94201 may represent our best example of a primitive basaltic melt (McSween *et al.* 1996), derived from the Martian interior, one can expect numerous experimental studies. Mikouchi *et al.* (2001), Koizumi *et al.* (2002), McKay *et al.* (2001) and McCanta and Rutherford (2001) have attempted to reproduce the pyroxenes found in QUE94201 under various $f\text{O}_2$ conditions. Karner *et al.* (2007, 2008) have

experimentally studied Cr and V partitioning between pyroxene and melt, and the valence state of V to exactly determine the reducing conditions of QUE94201.

Kring *et al.* (2003) review the “liquid line of descent” on the experimental phase diagrams (Longhi 1991). They note that the bulk composition falls within the olivine stability field rather than on the plagioclase pyroxene liquidus and offer various solutions to this “conundrum”. Experiments that include high phosphorus seem to be needed!

Weathering

Wentworth and Gooding (1996) have studied the weathering products in QUE94201. They report an abundance of Fe-sulfate, but since this is also observed in cavities in the fusion crust, this is almost certainly a weathering product of Antarctic origin (Harvey *et al.* 1996).

Processing

This small sample (12 g) had some remnant fusion crust which was difficult to distinguish from interior glass. The sample was initially thought to be a terrestrial rock, but the presence of maskelynite in thin section revealed its Martian origin (confirmed by Fe/Mn and oxygen isotope ratios). Allocations were made from small interior and exterior chips. Two potted butts were used to produce 13 thin sections.

QUE 94201 is listed as a “restricted” sample by the MWG (Score and Lindstrom 1993, page 5), because of its small size.

References for QUE94201

- Aramovich C.J., Herd Chris D.K. and Papike J.J. (2001) Possible causes for late-stage reaction textures associated with pyroxferroite and metastable pyroxenes in the basaltic Martian meteorites (abs#1003). *Lunar Planet. Sci.* **XXXII** Lunar Planetary Institute, Houston.
- Aramovich C.J., Herd C.D.K. and Papike J.J. (2002) Symplectites derived from metastable phases in Martian basaltic meteorites. *Amer. Mineral.* **87**, 1351-1359.
- Blichert-Toft Janne, Gleason J.D., Albarede F., Kring D.A., Hill D.H. and Boynton W.V. (1998a) The Hf isotopic compositions of Zagami and QUE94201: A garnet-free depleted Martian mantle (abs#1074). *Lunar Planet. Sci.* **XXIX** Lunar Planetary Institute, Houston.




Figure 7: X-ray map of whole thin section of QUE 94201. Pink is pyroxene, yellow is maskelynite (figure courtesy of Gordon McKay and Craig Schwandt).

Blichert-Toft J., Albarede F., Gleason J.D., Kring D.A., Hill D.H. and Boynton W.V. (1998b) Martian mantle evolution from the hafnium isotopic perspective (abs). *Meteorit. & Planet. Sci.* **33**, A16-17.

Blichert-Toft J., Gleason J.D., Telouk P. and Albarede F. (1999) The Lu-Hf isotope geochemistry of Shergottites and the evolution of the Martian mantle-crust system. *Earth Planet. Sci. Lett.* **173**, 25-39.

Bogard D.D. and Garrison D.H. (1999) Argon-39-argon-40 “ages” and trapped argon in Martian Shergottites,

Chassigny and Allan Hills 84001. *Meteorit. & Planet. Sci.* **34**, 451-473.

Borg Lars E., Nyquist L.E., Weismann H. and Shih ChiYu. (1996) Rb-Sr age and initial $^{87}\text{Sr}/^{86}\text{Sr}$ of basaltic Shergottite QUE94201(abs). *Meteorit. & Planet. Sci.* **31**, A18-A19.

Borg L.E., Nyquist Larry E., Taylor L.A., Wiesmann H. and Shih C-Y. (1997a) Rb-Sr and Sm-Nd isotopic analyses of QUE94201: Constraints on Martian differentiation processes (abs). *Lunar Planet. Sci.* **XXVIII**, 133-134. Lunar Planetary Institute, Houston.





Figure 8: X-ray maps of sector-zoned pyroxene in QUE94201 (from bottom left core of figure 7). (figure courtesy of Gordon McKay and Craig Schwandt).



- Borg L.E., Nyquist L.E., Taylor L.A., Wiesmann H. and Shih C-Y. (1997b) Constraints on Martian differentiation processes from Rb-Sr and Sm-Nd isotopic analyses of the basaltic Shergottite QUE94201. *Geochim. Cosmochim. Acta* **61**, 4915-4931.
- Clayton R.N. and Mayeda T.K. (1996) Oxygen isotopic studies of achondrites. *Geochim. Cosmochim. Acta* **60**, 1999-2017.
- Delaney J.S. and Dyar M.D. (2002a) Compositional and oxidation state zoning in Martian pyroxene: Paradox or process indicator (abs#1659). *Lunar Planet. Sci. XXXIII*. Lunar Planetary Institute, Houston.
- Dreibus G., Spettel B., Wlotzka F., Schultz L., Weber H.W., Jochum K.P. and Wänke H. (1996b) QUE94201: An unusual Martian basalt (abs). *Meteorit. & Planet. Sci.* **31**, A39-40.
- Dreibus G., Wänke H. and Lugmair G.W. (1997) Volatile inventories of Mars and Earth and their implications for the evolution of the planetary atmospheres (abs). In *Conference on Early Mars: Geologic and hydrologic evolution, physical and chemical environments, and the implications for life*. (eds. Clifford et al.) *LPI Contribution* **916**, 26. Lunar Planetary Institute, Houston.
- Eugster O., Weigel A. and Polnau E. (1996) Two different ejection events for basaltic Shergottites QUE94201, Zagami and Shergotty (2.6 Ma ago) and Lherzolithic Shergottites LEW88516 and ALHA77005 (3.5 Ma ago) (abs). *Lunar Planet. Sci. XXVII*, 345-346. Lunar Planetary Institute, Houston.
- Eugster O., Busemann H. and Lorenzetti S. (2002a) The pre-atmospheric size of Martian meteorites (abs#1096). *Lunar Planet. Sci. XXXIII* Lunar Planetary Institute, Houston.
- Eugster O., Busemann H., Lorenzetti S. and Terribilini D. (2002b) Ejection ages from ⁸¹Kr-⁸³Kr dating and pre-atmospheric sizes of Martian meteorites. *Meteorit. & Planet. Sci.* **37**, 1345-1360.
- Franchi I. A., Sexton A. S., Wright I.P. and Pillinger C.T. (1997) A refinement of oxygen isotopic composition of Mars (abs). *Lunar Planet. Sci. XXVIII*, 379-380. Lunar Planetary Institute, Houston.
- Gaffney Amy M., Borg Lars E. and Connelly J.N. (2006) U-Pb isotopic systematics of QUE94201: Seeing through terrestrial Pb contamination to identify and even lower-Mu source on Mars (abs#1483). *Lunar Planet. Sci. XXXVII* Lunar Planetary Institute, Houston.
- Gaffney A.M., Borg L.E. and Connelly J.N. (2007) Uranium-lead isotope systematics of Mars inferred from the basaltic Shergottite QUE 94201. *Geochim. Cosmochim. Acta* **71**, 5016-5031.
- Garrison D.H. and Bogard D.D. (1997) Argon-39/Argon-40 dating of Martian meteorites (abs). *Meteorit. & Planet. Sci.* **32**, A45.
- Garrison D.H. and Bogard D.D. (1998) Isotopic composition of trapped and cosmogenic noble gases in several Martian meteorites. *Meteorit. & Planet. Sci.* **33**, 721-736.
- Grady M.M., Verchovsky A., Wright I.P. and Pillinger C.T. (1996b) Fully automated stepped combustion analysis of carbon, nitrogen and Argon in QUE94201: Comparison with other Shergottites (abs). *Meteorit. & Planet. Sci.* **31**, A52-A53.
- Greenwood J.P., Riciputi L.R. and McSween H.Y. (1997b) Sulfide isotopic compositions in Shergottites and ALH84001 and possible implications for life on Mars. *Geochim. Cosmochim. Acta* **61**, 4449-4453.
- Greenwood J.P., Itoh S., Sakamoto N., Vicenzi E.P. and Yurimoto H. (2007a) D/H zoning in apatite of Martian meteorites QUE94201 and Los Angeles: Implications for water on Mars (abs#5437). *Meteorit. & Planet. Sci.* **45**, A68.
- Grossman J.N. and Score R. (1996) Recently classified specimens in the US Antarctic meteorite collection (1994-1996). *Meteorit. & Planet. Sci.* **31**, A161-A180.
- Harvey R.P., McCoy T.J. and Leshin L.A. (1996) Shergottite QUE94201: Texture, mineral compositions, and comparison with other basaltic Shergottites (abs). *Lunar Planet. Sci. XXVII*, 497-498. Lunar Planetary Institute, Houston.
- Herd C.D.K. and Papike J.J. (2000c) Oxygen fugacity of the Martian basalts from analysis of iron-titanium oxides: Implications for Mantle-crust interaction on Mars (abs). *Meteorit. & Planet. Sci.* **35**, A70.
- Herd C.D.K., Papike J.J. and Brearley A.J. (2001d) Oxygen fugacity of Martian basalts from electron microprobe oxygen and TEM-EELS analysis of Fe-Ti oxides. *Amer. Mineral.* **86**, 1015-1024.
- Karner J.M., Papike J.J., Sutton S.R., Shearer C.K., McKay G., Le L. and Burger P. (2007a) Valence state partitioning of Cr between pyroxene-melt: Effects of pyroxene and melt composition and direct determination of Cr valence states by XANES. Application to Martian basalt QUE94201 composition. *Amer. Mineral.* **92**, 2002-2005.

- Karner J.M., Papike J.J., Shearer C.K., McKay G., Le L. and Burger P. (2007b) Valance state partitioning of Cr and V between pyroxene-melt: Estimates of oxygen fugacity for Martian basalt QUE 94201. *Amer. Mineral.* **92**, 1238-1241.
- Karner J.M., Papike J.J., Sutton S.R., Schearer C.K., Burger P., McKay G. and Le L. (2008a) Valance state partitioning of V between pyroxene-melt: Effects of pyroxene and melt composition, and direct determination of V valance states by XANES. Application to Martian basalt QUE 94201 composition. *Meteorit. & Planet. Sci.* **43**, 1275-1285.
- Karner J.M., Papike J.J., McKay G., Sutton S.R., Schearer C.K., Burger P. and Le L. (2008b) The partitioning of Cr and V between pyroxene-melt in Martian basalt QUE 94201 (abs#5254). *Meteorit. & Planet. Sci.* **43**, A70.
- Koizumi E., Mikouchi T., McKay G.A., Le L. and Schwandt C. (2001) Dynamic crystallization of the Queen Alexandra Range 94201 Shergottite (abs). *Meteorit. & Planet. Sci.* **36**, A102.
- Koizumi E., McKay G.A., Mikouchi T., Le L., Schwandt C., Monkawa A. and Miyamoto M. (2002a) Crystallization experiments of the Martian meteorite QUE94201: Additional constraints on its formation condition (abs#1442). *Lunar Planet. Sci.* **XXXIII** Lunar Planetary Institute, Houston.
- Koizumi E., Mikouchi T., McKay G.A., Le L., Schwandt C., Monkawa A. and Miyamoto M. (2002b) Comparative cooling experiments of Queen Alexandra Range 94201 basaltic Shergottite under reducing and oxidizing conditions (abs). *Meteorit. & Planet. Sci.* **37**, A80.
- Koizumi E., Mikouchi T., McKay G., Le L., Schwandt C., Monkawa A. and Miyamoto M. (2002c) Effects of cooling rate and oxygen fugacity on the crystallization of the Queen Alexander Range 94201 Martian melt composition (abs). *Antarctic Meteorites XXVII*, 63-65. Nat. Inst. Polar Res., Tokyo.
- Koizumi E., Mikouchi T., McKay G. and Miyamoto M. (2008) Crystallization relationship of Yamato 980459 and QUE94201 depleted Shergottites (abs#5127). *Meteorit. & Planet. Sci.* **43**, A79.
- Kring D.A. (2002) QUE 94201: Reconsidering its origins as a bulk melt from a volcanic region of Mars (abs). Unmixing SNCs. 31-32. LPI Contribution No. 1134. Lunar Planetary Institute, Houston.
- Kring D.A., Gleason J.D., Hill D.H., Jull A.J.T. and Boynton W.V. (1996) QUE94201, a new Martian meteorite that may represent a bulk melt rather than a cumulate fraction (abs). *Lunar Planet. Sci.* **XXVII**, 705-706. Lunar Planetary Institute, Houston.
- Kring D.A., Gleason J.D., Swindle T.D., Nishiizumi K., Caffee M.W., Hill D.H., Jull A.J.T. and Boynton W.V. (2003) Composition of the first bulk melt sample from a volcanic region on Mars: Queen Alexandra Range 94201. *Meteorit. & Planet. Sci.* **38**, 1833-1848.
- Lentz R.C.F. and McSween H.Y. (1999b) Basaltic Shergottite crystallization: A quantitative textural analysis (abs). *Meteorit. & Planet. Sci.* **34**, A74.
- Lentz R.C.F. and McSween H.Y. (2000) Crystallization of the basaltic Shergottites: Insights from crystal size distribution (CSD) analysis of pyroxenes. *Meteorit. & Planet. Sci.* **35**, 919-927.
- Lentz R.C.F., McSween H.Y. and Fayek M. (2004) Light lithophile abundances and isotopic ratios in Shergottites (abs#1623). *Lunar Planet. Sci.* **XXXV** Lunar Planetary Institute, Houston.
- Leshin L.A. (1998a) The origin and evolution of water on Mars: Possible constraints on the Juvenile reservoir (abs#1468). *Lunar Planet. Sci.* **XXIX** Lunar Planetary Institute, Houston.
- Leshin L.A. (2000) Insight into martian water reservoirs from analyses of Martian meteorite QUE94201. *Geophys. Res. Lett.* **27**, 2017-2020.
- Leshin L.A., Harvey R.P., McCoy T.J. and McKeegan K.D. (1996b) Water in apatite from Shergottite QUE94201: Abundances and D/H (abs). *Meteorit. & Planet. Sci.* **31**, A79-A80.
- Leshin L.A., Epstein S. and Stolper E.M. (1996c) Hydrogen isotope geochemistry of SNC meteorites. *Geochim. Cosmochim. Acta* **60**, 2635-2650.
- McCanta Molly and Rutherford M.J. (2001) SNC oxygen fugacity as recorded in pyroxenes: An experimental study (abs#1348). *Lunar Planet. Sci.* **XXXII** Lunar Planetary Institute, Houston.
- McKay G.A., Yang S.-R. and Wagstaff J. (1996) Complex zoned pyroxenes in Shergottite QUE94201: Evidence for a two-stage crystallization history (abs). *Lunar Planet. Sci.* **XXVII**, 851-852. Lunar Planetary Institute, Houston.
- McKay G.A., Koizumi E., Mikouchi T., Le L. and Schwandt C. (2001) Experimental crystallization of the QUE94201 basaltic Shergottite: Support for the Martian magma hypothesis (abs). *NIPR Sym. Antarctic Meteorites 24th*, 77-79. Nat. Inst. Polar Res., Tokyo.

- McKay G.A., Koizumi E., Mikouchi T., Le L. and Schwandt C. (2002a) Crystallization of Shergottite QUE94201: An experimental study (abs#2051). *Lunar Planet. Sci.* **XXXIII** Lunar Planetary Institute, Houston.
- McKay G.A., Le L., Koizumi E. and Mikouchi T. (2003) Additional constraints on the crystallization of basaltic Shergottite QUE94201 (abs#2109). *Lunar Planet. Sci.* **XXXIV** Lunar Planetary Institute, Houston.
- McSween H.Y. and Eisenhour D.D. (1996) QUE94201, A noncumulate Shergottite (abs). *Lunar Planet. Sci.* **XXVII**, 853-854. Lunar Planetary Institute, Houston.
- McSween H.Y., Eisenhour D.D., Taylor L.A., Wadhwa M. and Crozaz G. (1996) QUE94201 Shergottite: Crystallization of a Martian basaltic magma. *Geochim. Cosmochim. Acta* **60**, 4563-4569.
- Mikouchi T. (2000) Pyroxene and plagioclase in the Los Angeles Martian meteorite: Comparison with the Queen Alexandra Range 94201 Martian meteorite and the Asuka 881757 lunar meteorite (abs). *Meteorit. & Planet. Sci.* **35**, A110.
- Mikouchi T., Miyamoto M. and McKay G.A. (1996) Mineralogy and petrology of new Antarctic Shergottite QUE94201: A coarse-grained basalt with unusual pyroxene zoning (abs). *Lunar Planet. Sci.* **XXVII**, 879-880. Lunar Planetary Institute, Houston.
- Mikouchi T., Miyamoto M. and McKay G. A. (1997a) Similarities in zoning of pyroxenes from QUE94201 and EETA79001 Martian meteorites (abs). *Lunar Planet. Sci.* **XXVIII**, 955-956. Lunar Planetary Institute, Houston.
- Mikouchi T., Miyamoto M. and McKay G. A. (1997b) Crystallization histories of basaltic Shergottites as inferred from chemical zoning of pyroxene and maskelynite (abs). *Meteorit. & Planet. Sci.* **32**, A92-93.
- Mikouchi T., Miyamoto M. and McKay G.A. (1998a) Mineralogy of Antarctic basaltic Shergottite Queen Alexandra Range 94201: Similarities to Elephant Moraine A79001 (Lithology B) Martian meteorite. *Meteorit. & Planet. Sci.* **33**, 181-189.
- Mikouchi T., Osaka T., Kaneda K. and Ohsumi K. (1998d) X-ray diffraction study of shocked plagioclase in Martian and Lunar meteorites with the micro-area Laue method using synchrotron radiation (abs). *NIPR Sym. Antarctic Meteorites* **23rd**, 80-82. Nat. Inst. Polar Res., Tokyo.
- Mikouchi T., Miyamoto M. and McKay G.A. (1999d) The role of undercooling in producing igneous zoning trends in pyroxenes and maskelynites among basaltic Martian meteorites. *Earth Planet. Sci. Lett.* **173**, 235-256.
- Mikouchi T., Koizumi E., McKay G., Le L. and Schwandt C. (2001a) Experimental crystallization of the QUE94201 basaltic Shergottite (abs#2100). *Lunar Planet. Sci.* **XXXII** Lunar Planetary Institute, Houston.
- Mikouchi T. and McKay G. (2003) Shock heating and subsequent cooling of basaltic Shergottites: The case for QUE94201 and Dhofar 378 (abs#1920). *Lunar Planet. Sci.* **XXXIV**. Lunar Planetary Institute, Houston.
- Mittlefehldt D.W. and Lindstrom M.M. (1996) Martian meteorites QUE94201, an unusual basalt, and Governador Valadares, a typical clinopyroxenite: Geochemistry (abs). *Lunar Planet. Sci.* **XXVII**, 887-888. Lunar Planetary Institute, Houston.
- Mohapatra R.K., Schwenzer S.P., Herrmann S. and Ott U. (2002b) Nitrogen and Noble gases in the basaltic Shergottite QUE94201 (abs). *Meteorit. & Planet. Sci.* **37**, A102.
- Nishiizumi K. and Caffee M.W. (1996) Exposure history of Shergottite Queen Alexandra Range 94201 (abs). *Lunar Planet. Sci.* **XXVII**, 961-962. Lunar Planetary Institute, Houston.
- Papike J.J., Karner J.M. and Shearer C.K. (2003a) Determination of planetary basalt parentage: A simple technique using the electron microprobe (abs#1018). *Lunar Planet. Sci.* **XXXIV** Lunar Planetary Institute, Houston.
- Papike J.J., Karner J.M. and Shearer C.K. (2003b) Determination of planetary basalt parentage: A simple technique using the electron microprobe. *Amer. Mineral.* **88**, 469-472.
- Papike J.J., Karner J.M. and Shearer C.K. (2005) Comparative planetary mineralogy: Valence state partitioning of Cr, Fe, Ti, and V among crystallographic sites in olivine, pyroxene and spinel from planetary basalts. *Amer. Mineral.* **90**, 277-290.
- Papike J.J., Karner J.M., Shearer C.K. and Burger P.V. (2009a) Pyroxene mineralogy of Martian meteorites (abs#1180). *Lunar Planet. Sci.* **XL**, Lunar Planetary Institute, The Woodlands.
- Papike J.J., Karner J.M., Spilde M.N., Shearer C.K. and Burger P.V. (2009b) Silicate mineralogy of Martian meteorites. *Geochim. Cosmochim. Acta* **73**, 7443-7485. (invited review with great pictures of textures)
- Papike J.J., Burger P.V., Shearer C.K., McCubbin F.M. and Elardo S.M. (2012) Experimental Martian eclogite with a QUE94201 composition (abs#1010). *Lunar Planet. Sci.* **XLIII**, Lunar Planetary Institute, The Woodlands.

- Schnabel C., Ma. P., Herzog G.F., Faestermann T., Knie K. and Korschinek G. (2001) ^{10}Be , ^{26}Al and ^{53}Mn in Martian meteorites (abs#1353). *Lunar Planet. Sci.* **XXXII**, Lunar Planetary Institute, Houston.
- Score R. and Mason B. (1995) Macroscopic and thin section description of QUE94201. In *Antarctic Meteorite Newsletter* **18 (2)**, 20. JSC Curator's Office, Houston.
- Simon Justin I., DePaolo D.J. and Moynier F. (2009) Calcium isotope composition of meteorites, Earth and Mars. *The Astrophys. Jour.* **702**, 707-715.
- Swindle T.D., Li B. and Kring D.A. (1996) Noble gases in Martian meteorite QUE94201 (abs). *Lunar Planet. Sci.* **XXVII**, 1297-1298. Lunar Planetary Institute, Houston.
- Papike J.J., Karner J.M., Spilde M.N., Shearer C.K. and Burger P.V. (2009b) Silicate mineralogy of Martian meteorites. *Geochim. Cosmochim. Acta* **73**, 7443-7485. (invited review with great pictures of textures)
- Terribilini D., Busemann H. and Eugster O. (2000) Krypton-81-Krypton cosmic-ray exposure ages of the Martian meteorites including the new Shergottite Los Angeles (abs). *Meteorit. & Planet. Sci.* **35**, A155-156.
- Wadhwa M. and Crozaz G. (1996) QUE94201: A new and different Shergottite (abs). *Lunar Planet. Sci.* **XXVII**, 1365-1366. Lunar Planetary Institute, Houston.
- Wadhwa M., Crozaz G., Taylor L.A., McSween H. Y. (1997a) Martian basalt (Shergottite) QUE94201 and lunar basalt 15555: A tale of two pyroxenes (abs). *Lunar Planet. Sci.* **XXVIII**, 1485-1486. Lunar Planetary Institute, Houston.
- Wadhwa M., Crozaz G., Taylor L.A. and McSween H.Y. (1998a) Martian basalt (Shergottite) Queen Alexandra Range 94201 and lunar basalt 15555; A tale of two pyroxenes. *Meteorit. & Planet. Sci.* **33**, 321-328.
- Wang M-S., Mokoš J. A. and Lipschutz M. E. (1999) Martian meteorites: Volatile trace elements and cluster analysis. *Meteorit. & Planet. Sci.* **33**, 671-675.
- Warren P.H., Kallemeyn G.W., Arai T. and Kaneda K. (1996) Compositional-petrologic investigations of Eucrites and the QUE94201 Shergottite (abs). *Proc. NIPR Sym. Antarctic Meteorites* **21st**, 195-197. Nat. Inst. Polar Res., Tokyo.
- Warren P.H. and Kallemeyn G.W. (1997a) Yamato-793605 and other presumed Martian meteorites: Composition and petrogenesis (abs). *NIPR Sym. Antarctic Meteorites* **22nd**, 197-199. Nat. Inst. Polar Res., Tokyo.
- Warren P.H. and Kallemeyn G.W. (1997b) Yamato-793605, EET79001 and other presumed Martian meteorites: Compositional clues to their origins. *Antarctic Meteorite Research* **10**, 61-81. Nat. Inst. Polar Res., Tokyo.
- Wentworth S.J. and Gooding J.L. (1996) Water-based alteration of the Martian meteorite, QUE94201, by sulfate-dominated solutions (abs). *Lunar Planet. Sci.* **XXVII**, 1421-1422. Lunar Planetary Institute, Houston.