

15065
Gabbroic Basalt
1475.5 grams




Figure 1: Photo of 15065,13 set up in wire saw apparatus. Cube is 1 cm. NASA S71-60957.

Introduction

15065 is a coarse-grained mare basalt that is about 3.35 b.y. old. It was collected as part of a suite of samples at different distances from the rim of Elbow Crater (a radial sample collection). Indeed it was the coarsest basalt (of the sequence 15065, 15075 and 15485) and closest to the crater, perhaps from the deepest layer of the basalt flow (Swann et al. 1971).

15065 may not be related to either the olivine-normative, nor quartz-normative suites of Apollo 15 basalts (Ryder 1985). In any case the vuggy, mafic portion of 15065 deserves more study.

Petrography

Ryder (1985) termed 15065 a coarse-grained quartz-normative basalt. However, it has two distinct regions with different feldspar to mafic mineral content (originally noted by Morrison and Silver in the original Apollo 15 catalog (Butler 1972). 15065 has abundant prismatic/euhedral pigeonite crystals with greenish cores and red-brown rims intergrown with anhedral to subhedral plagioclase (figures 1, 2 and 3). Ragged anhedral olivine grains (Fo₅₀) are found enclosed in pigeonite cores.

Mineralogical Mode for 15065

	Sample Catalog Butler 1971	Brown et al. 1972	Longhi et al. 1972	Juan et al. 1972
Olivine		2	1.3	
Pyroxene	60		63	70
Plagioclase	35		31.6	27
Opaque	1		2.2	2
Silica	1		1.9	



Figure 2: PET photo of 15065 showing mafic mineral segregation on right side near the 1 inch cube. NASA S71-42924.




Figure 3: Photomicrograph of thin section 15065,89 by C Meyer @ 20x.

Brown et al. (1972), Gay et al. (1972) and others recognized that 15065 was one of the more slowly cooled basalts. Lofgren et al. (1975), Grove and Walker (1977), Taylor et al. (1976) and Onorato et al. (1979) performed mineral analyses and laboratory experiments to determine the cooling rate.

Mineralogy

Olivine: Mg-rich olivine is found as cores in pigeonite, while fayalite is found in the interstitial residuum.

Pyroxene: The chemical composition of pyroxene in 15065 was reported by Grove and Bence (1977)(figure 4). Yajima and Hafner (1974) determined the equilibrium temperature from the cation distribution in pigeonite.

Plagioclase: Longhi et al. (1972, 1976) studied the Mg and Fe content of plagioclase (An_{80-91}).

Ilmenite: The Zr content of ilmenite has been reported (Taylor and McCallister 1972, Blank et al. 1982).

Chromite: Taylor et al. (1975) and El Goresy et al. (1976) studied chromite and ulvospinel.

Metal: Taylor et al. (1975) found high Co content of metal grains in the whole suite of samples from Elbow




Figure 4: Pyroxene and olivine composition of 15065.

Crater (figure 5). Jedwab (1972) described faceted, isometric iron grains from vugs.

Phosphate: Prominent accessory apatite is found intergrown with ferro-augite, sulfides and silica (both tridymite and cristobalite reported).

Zirconolite: Wark et al. (1973) reported the composition of zirconolite (determined by ElGoresy).

Chemistry

Nava (1975) found that the two different regions in 15065 were indeed different in composition (tables 1 and 2). The more mafic region has high Fe and Ti (figure 7). Wanke and others reported the trace element content (figure 6). The composition of 15065 can't be related to either the quartz-normative, nor olivine-normative suites of Apollo 15 basalts by simple olivine removal.

Gibson et al. (1975) determined the sulfur content (600 ppm).

Barker (1974) and Gibson and Andrawes (1978) attempted to determine the composition of gases released by heating 15065 (figure 10).

Radiogenic age dating

Papanastassiou and Wasserburg (1973) and Nakamura et al. (1977) determined the age of 15065 (figures 8 and 9). Unruh et al. (1984) reported the isotopic composition of Lu-Hf and Sm-Nd.

Cosmogenic isotopes and exposure ages

Eldridge et al. (1972) determined the cosmic-ray exposure of $^{22}\text{Na} = 23 \text{ dpm/kg}$, $^{26}\text{Al} = 68 \text{ dpm/kg}$ and




Figure 5: Composition of metal grains in basalt samples from Elbow Crater (from Taylor et al. 1975).




Figure 6: Trace element analysis of 15565 compared with that of 15601.




Figure 7: Composition of 15065 compared with that of other lunar basalts.




Figure 8: Rb/Sr isochrons for various Apollo 15 basalts including 15065 (top). This is reproduced from Papanastassiou and Wasserburg (1973).

$^{54}\text{Mn} = 27 \text{ dpm/kg}$, but the cosmic ray exposure age hasn't been reported.

Other Studies

Hargraves and Dorety (1972) studied the magnetic properties, Adams and McCord (1972) and Charrette and Adams (1975) used 15065 to obtain reflectance spectra, Chung and Westphal (1973) studied the physical properties and Bhattacharya et al. (1973) studied tracks.

Processing

A portion of the rather vuggy, mafic portion of 15065 was separated from the main mass of 15065 before sawing. Additional mafic material can be found on piece (.117). Figure 1 shows that ,13 was cut with the wire saw (a rather messy proposition).

There are 18 thin sections of 15065. Sections number ,87 and ,91 are from the mafic portion.




Figure 9: Sm/Nd isochron for 15565 (Nakamura et al. 1977).

Summary of Age Data for 15065

	Rb/Sr	Sm/Nd
Papanastassiou and Wasserburg 1973	$3.28 \pm 0.04 \text{ b.y.}$	
Nakamura et al. 1977		3.34 ± 0.09

Caution: These are calculated with original decay constants !




Figure 10: Gas release curves for thermal heating of 15065 (Gibson and Moore 1972).

Table 1a. Chemical composition of 15065 (main portion).

reference weight	Cuttitta73 Christian72	Juan 72	O'Kelley72	Strasheim72	Wanke 75	Nava 75	Fruchter73	Ganapathy73 ,5	Baedecker73
SiO2 %	48.47	(a) 43.2	(b)	47.24	(d) 48.99	(f) 48.2	(g)		
TiO2	1.48	(a) 1.88	(b)	1.54	(d) 1.33	(f) 1.44	(g)		
Al2O3	9.26	(a) 10.1	(b)	9.33	(d) 12.75	(f) 10.32	(g) 11.34	(f)	
FeO	19.18	(a) 19.2	(b)	19.17	(d) 17.75	(f) 18.46	(g) 19.17	(f)	
MnO	0.26	(a) 0.3	(b)	0.26	(d) 0.24	(f) 0.234	(g) 1.22	(f)	
MgO	10.57	(a) 11.5	(b)	10.69	(d) 8.8	(f) 10.35	(g)		
CaO	9.94	(a) 12.12	(b)	9.53	(d) 11.23	(f) 9.55	(g)		
Na2O	0.34	(a) 0.715	(b)	0.23	(d) 0.37	(f) 0.33	(g) 0.28	(f)	
K2O	0.05	(a) 0.068	(b) 0.046	(c) 0.05	(d) 0.05	(f) 0.041	(g)		
P2O5	0.05	(a)		0.08	(d) 0.06	(f) 0.104	(g)		
S %					0.05	(f)			
sum									
Sc ppm	38	(a)		88	(d) 39.1	(f)	41	(f)	
V	158	(a)		185	(d)				
Cr	3626	(a)		4379	(d) 3160	(f)	4600	(f)	
Co	52	(a) 70		47	(d) 37.7	(f)	46	(f) 45	(f)
Ni	151	(a) 147		78	(d)	(f)			63 (f)
Cu	64	(a) 56		48	(d) 5.42	(f)			
Zn		38			<1	(f)	1.6	(f) 0.92	(f)
Ga	4.1	(a) 24			3.76	(f)			3.1 (f)
Ge ppb					<100	(f)	5.3	(f) 21	(f)
As					0.9	(f)			
Se					0.08	(f)	167	(f)	
Rb	<1	(a)		5	(d) 0.7	(f)	0.76	(f)	
Sr	110	(a) 214		98	(d) 134	(f)			
Y	23	(a)		29	(d) 24	(f)			
Zr	63	(a)		79	(d) 94	(f)			
Nb	12	(a)		13	(d) 6	(f)			
Mo									
Ru									
Rh									
Pd ppb									
Ag ppb		0.545					0.91	(f)	
Cd ppb							1.5	(f) 1.2	(f)
In ppb							0.51	(f) 0.32	(f)
Sn ppb									
Sb ppb							0.016	(f)	
Te ppb							3.4	(f)	
Cs ppm					0.05	(f)	0.062	(f)	
Ba	50	(a)		85	(e) 96	(f)			
La	<10	(a)		7.1	(e) 7.73	(f)	5.1	(f)	
Ce				30	(e) 20.6	(f)			
Pr				3.7	(e) 3.15	(f)			
Nd				20	(e) 14.6	(f)			
Sm				5	(e) 4.72	(f)	3.4	(f)	
Eu				1.05	(e) 1.14	(f)	0.94	(f)	
Gd				5.6	(e) 5.3	(f)			
Tb				0.86	(e) 0.96	(f)	0.7	(f)	
Dy				6.8	(e) 6.66	(f)	2	(f)	
Ho				1.25	(e) 1.2	(f)			
Er				3.9	(e) 3.7	(f)			
Tm				0.47	(e)				
Yb	3.8			2.9	(e) 2.98	(f)	2	(f)	
Lu				0.3	(e) 0.43	(f)	0.34	(f)	
Hf					3.36	(f)	2.1	(f)	
Ta					0.45	(f)	0.4	(f)	
W ppb					102	(f)			
Re ppb					<0.1	(f)			
Os ppb							0.0015	(f)	
Ir ppb							0.0054	(f) 0.048	(f)
Pt ppb									
Au ppb		4			0.031	(f)			
Th ppm			0.51	(c)	0.7	(f)	0.021	(f) 0.19	(f)
U ppm			0.15	(c)	0.19	(f)	0.235	(f)	

technique: (a) conventional, (b) ?, (c) radiation counting, (d) ave. of mixed, (e) MS, (f) INAA, RNAA


Table 1b. Chemical composition of 15065 (mafic portion).

reference weight	Nava 75	Cuttitta73 Christian72	Ganapathy73 ,41	
SiO2 %	47.7	(g) 47.95	(a)	
TiO2	2.86	(g) 2.3	(a)	
Al2O3	6.05	(g) 5.33	(a)	
FeO	23.77	(g) 23.6	(a)	
MnO	0.307	(g) 0.31	(a)	
MgO	9.52	(g) 10.15	(a)	
CaO	9.33	(g) 9.3	(a)	
Na2O	0.27	(g) 0.25	(a)	
K2O	0.081	(g) 0.07	(a)	
P2O5	0.119	(g) 0.09	(a)	
S %				
sum				
Sc ppm		53	(a)	
V		178	(a)	
Cr		3421	(a)	
Co		66	(a)	40 (f)
Ni		54	(a)	
Cu		14	(a)	
Zn				1 (f)
Ga		5.3	(a)	
Ge ppb				6.7 (f)
As				
Se			53	(f)
Rb		1	(a)	0.39 (f)
Sr		100	(a)	
Y		39	(a)	
Zr		103	(a)	
Nb		<10	(a)	
Mo				
Ru				
Rh				
Pd ppb				
Ag ppb				0.88 (f)
Cd ppb				1.1 (f)
In ppb				0.18 (f)
Sn ppb				
Sb ppb				0.24 (f)
Te ppb				0.96 (f)
Cs ppm				0.026 (f)
Ba		75	(a)	
La		22	(a)	
Ce				
Pr				
Nd				
Sm				
Eu				
Gd				
Tb				
Dy				
Ho				
Er				
Tm				
Yb		6.6		
Lu				
Hf				
Ta				
W ppb				
Re ppb				0.0094 (f)
Os ppb				
Ir ppb				0.144 (f)
Pt ppb				
Au ppb				0.014 (f)
Th ppm				
U ppm				0.085 (f)

technique: (a) conventional, (b) ?, (c) radiation counting, (d) ave. of mixed, (e) MS, (f) INAA, RNAA



Figure 11: Photo of 15065,117 showing area of mafic mineral segregation. NASA S74-27569. Sample is 11 cm across.



References for 15065

- Adams J.B. and McCord T.B. (1972) Optical evidence for average pyroxene composition of Apollo 15 samples. In **The Apollo 15 Lunar Samples**, 10-13. Lunar Planetary Institute, Houston.
- Baedecker P.A., Chou C.-L., Grudewicz E.B. and Wasson J.T. (1974) Volatile and siderophile trace elements in Apollo 15 samples: Geochemical implications and characterization of the long-lived and short-lived extralunar materials. *Proc. 4th Lunar Sci. Conf.* 1177-1195.
- Barker C. (1974) Composition of the gases associated with the magmas that produced rocks 15016 and 15065. *Proc. 5th Lunar Sci. Conf.* 1737-1746.
- Bhattacharya S.K., Goswami J.N., Lal D., Patel P.P. and Rao M.N. (1975) Lunar regolith and gas-rich meteorites: Characterization based on particle tracks and grain-size distributions. *Proc. 6th Lunar Sci. Conf.* 3509-3526.
- Bibring J.P., Burlingame A.L., Langevin Y., Maurette M. and Wszolek P.C. (1974) Simulation of lunar carbon chemistry: II. Solar winds contribution. *Proc. 5th Lunar Sci. Conf.* 1763-1784.
- Blank H., ElGoresy A., Janicke J., Nobile R. and Traxel K. (1984) Partitioning of Zr and Nb between coexisting opaque phases in lunar rocks - determined by quantitative proton microprobe analysis. *Earth Planet. Sci. Lett.* **68**, 19-33.
- Brown G.M., Emeleus C.H., Holland G.J., Peckett A. and Phillips R. (1972) Mineral-chemical variations in Apollo 14 and Apollo 15 basalts and granitic fractions. *Proc. 3rd Lunar Sci. Conf.* 141-157.
- Butler P. (1971) Lunar Sample Catalog, Apollo 15. Curators' Office, MSC 03209
- Carron M.K., Annel C.S., Christian R.P., Cuttitta F., Dwornik E.J., Ligon D.T. and Rose H.J. (1972) Elemental analysis of lunar soil samples from Apollo 15 mission. In **The Apollo 15 Samples** 198-201.
- Charette M.P. and Adams J.B. (1975a) Mare basalts: Characterization of compositional parameters by spectral reflectance. In *Papers presented to the Conference on Origins of Mare Basalts and their Implications for Lunar Evolution*, 25-28. Lunar Planetary Institute, Houston.
- Chung D.H. and Westphal W.B. (1973) Dielectric spectra of Apollo 15 and 16 lunar solid samples. *Proc. 4th Lunar Sci. Conf.* 3077-3091.
- Chung Dae H. (1973) Elastic wave velocities in anorthosite and anorthositic gabbros from Apollo 15 and 16 landing sites. *Proc. 3rd Lunar Sci. Conf.* 2591-2600.
- Clayton R.N., Hurd J.M. and Mayeda T.K. (1973) Oxygen isotope compositions of Apollo 15, 16 and 17 samples, and their bearing on lunar origin and petrogenesis. *Proc. 4th Lunar Sci. Conf.* 1535-1542.
- Cuttitta R., Rose H.J., Annel C.S., Carron M.K., Christian R.P., Ligon D.T., Dwornik E.J., Wright T.L. and Greenland L.P. (1973) Chemistry of twenty-one igneous rocks and soils returned by the Apollo 15 mission. *Proc. 4th Lunar Sci. Conf.* 1081-1096.
- Eldridge J.S., O'Kelley G.D. and Northcutt K.J. (1972) Concentrations of cosmogenic radionuclides in Apollo 15 rocks and soils. In **The Apollo 15 Lunar Samples** 357-359. Lunar Sci. Institute, Houston.
- El Goresy A., Prinz M. and Ramdohr P. (1976a) Zoning in spinels as an indicator of the crystallization histories of mare basalts. *Proc. 7th Lunar Sci. Conf.* 1261-1279.
- Fruchter J.S., Stoesser J.W., Lindstrom M.M. and Goles G.G. (1973) Apollo 15 clastic materials and their relationship to local geologic features. *Proc. 4th Lunar Sci. Conf.* 1227-1237.
- Ganapathy R., Morgan J.W., Krahenbuhl U. and Anders E. (1973) Ancient meteoritic components in lunar highland rocks: Clues from trace elements in Apollo 15 and 16 samples. *Proc. 4th Lunar Sci. Conf.* 1239-1261.
- Gay P., Muir I.D. and Price G.G. (1972) Mineralogy and petrology of two Apollo 15 mare basalts. In **The Apollo 15 Lunar Samples**. 70-72. (Ed. Chaimberlain and Watkins) LPI
- Gibson E.K. and Moore G.W. (1972c) Thermal analysis-inorganic gas release studies on Apollo 14, 15 and 16 lunar samples. In **The Apollo 15 Lunar Samples**. 307-310.
- Gibson E.K. and Andrawes F.F. (1978a) Nature of the gases released from lunar rocks and soils upon crushing. *Proc. 9th Lunar Planet. Sci. Conf.* 2433-2450.
- Gibson E.K., Chang S., Lennon K., Moore G.W. and Pearce G.W. (1975a) Sulfur abundances and distributions in mare basalts and their source magmas. *Proc. 6th Lunar Sci. Conf.* 1287-1301.
- Grove T.L. and Bence A.E. (1977) Experimental study of pyroxene-liquid interactions in quartz-normative basalt 15597. *Proc. 8th Lunar Sci. Conf.* 1549-1579.

- Hargraves R.B. and Dorety N. (1972b) Remanent magnetism in four Apollo 15 igneous rock fragments. *In **The Apollo 15 Lunar Samples*** 415-417. Lunar Planetary Institute, Houston.
- Jedwab J. (1972) Mineralogical notes on Apollo 15 samples. *In **The Apollo 15 Lunar Samples***, 108-109.
- Juan V.C., Chen J.C., Huang C.K., Chen P.Y. and Wang Lee C.M. (1972b) Petrology and chemistry of some Apollo 15 crystalline rocks. *In **The Apollo 15 Lunar Samples***, 110-115.
- Lofgren G.E., Donaldson C.H. and Usselman T.M. (1975) Geology, petrology and crystallization of Apollo 15 quartz-normative basalts. *Proc. 6th Lunar Sci. Conf.* 79-99.
- Longhi J., Walker D., Stolper E.N., Grove T.L. and Hays J.F. (1972) Petrology of mare/rille basalts 15555 and 15065. *In **The Apollo 15 Lunar Samples***, 131-134. Lunar Sci. Institute, Houston.
- Longhi J., Walker D. and Hays J.F. (1976) Fe and Mg in plagioclase. *Proc. 7th Lunar Sci. Conf.* 1281-1300.
- LSPET (1972a) The Apollo 15 lunar samples: A preliminary description. *Science* **175**, 363-375.
- LSPET (1972b) Preliminary examination of lunar samples. Apollo 15 Preliminary Science Report. NASA SP-289, 6-1—6-28.
- Moore C.B., Lewis C.F., and Gibson E.K. (1972) Carbon and nitrogen in Apollo 15 lunar samples. *In **The Apollo 15 Lunar Samples*** (Chamberlain and Watkins, eds.), 316-318. The Lunar Science Institute, Houston.
- Moore C.B., Lewis C.F. and Gibson E.K. (1973) Total carbon contents of Apollo 15 and 16 lunar samples. *Proc. 4th Lunar Sci. Conf.* 1613-1923.
- Nakamura N., Unruh D.M., Gensho R. and Tatsumoto M. (1977) Evolution history of lunar mare basalts: Apollo 15 samples revisited (abs). *Lunar Sci.* **VIII**, 712-714.
- Nava D.F. (1974a) Chemical compositions of some soils and rock types from the Apollo 15, 16, and 17 lunar sites. *Proc. 5th Lunar Sci. Conf.* 1087-1096.
- O'Kelley G.D., Eldridge J.S. and Northcutt K.J. (1972a) Abundances of primordial radioelements K, Th, and U in Apollo 15 samples, as determined by non-destructive gamma-ray spectrometry. *In **The Apollo 15 Lunar Samples*** (Chamberlain J.W. and Watkins C., eds.), 244-246. Lunar Science Institute, Houston.
- O'Kelley G.D., Eldridge J.S., Schonfeld E. and Northcutt K.J. (1972b) Primordial radionuclides and cosmogenic radionuclides in lunar samples from Apollo 15. *Science* **175**, 440-443.
- O'Kelley G.D., Eldridge J.S., Northcutt K.J. and Schonfeld E. (1972c) Primordial radionuclides and cosmogenic radionuclides in lunar samples from Apollo 15. *Proc. 3rd Lunar Sci. Conf.* 1659-1670.
- Onorato P.I.K., Yinnon H., Uhlmann D.R. and Taylor L.A. (1979) Partitioning as a cooling rate indicator. *Proc. 10th Lunar Planet. Sci. Conf.* 479-491.
- Papanastassiou D.A. and Wasserburg G.J. (1973) Rb-Sr ages and initial strontium in basalts from Apollo 15. *Earth Planet. Sci. Lett.* **17**, 324-337.
- Papike J.J., Hodges F.N., Bence A.E., Cameron M. and Rhodes J.M. (1976) Mare basalts: Crystal chemistry, mineralogy and petrology. *Rev. Geophys. Space Phys.* **14**, 475-540.
- Ryder G. (1985) Catalog of Apollo 15 Rocks (three volumes). Curatorial Branch Pub. # 72, JSC#20787
- Strasheim A., Coetzee J.H.J., Jackson P.F.S., Strelow F.W.E., Wybenga F.T., Gricius A.J. and Kokot M.L. (1972b) Analysis of lunar samples 15065, 15301, and 15556, with isotopic data for ⁷Li/⁶Li. *In **The Apollo 15 Lunar Samples*** (Chamberlain and Watkins, eds.), 257-259. The Lunar Science Institute, Houston.
- Swann G.A., Hait M.H., Schaber G.C., Freeman V.L., Ulrich G.E., Wolfe E.W., Reed V.S. and Sutton R.L. (1971b) Preliminary description of Apollo 15 sample environments. U.S.G.S. Interagency report: 36. pp219 with maps
- Swann G.A., Bailey N.G., Batson R.M., Freeman V.L., Hait M.H., Head J.W., Holt H.E., Howard K.A., Irwin J.B., Larson K.B., Muehlberger W.R., Reed V.S., Rennilson J.J., Schaber G.G., Scott D.R., Silver L.T., Sutton R.L., Ulrich G.E., Wilshire H.G. and Wolfe E.W. (1972) 5. Preliminary Geologic Investigation of the Apollo 15 landing site. In Apollo 15 Preliminary Science Rpt. NASA SP-289. pages 5-1-112.
- Tatsumoto M., Hedge C.E., Knight R.J., Unruh D.M. and Doe Bruce R. (1972b) U-Th-Pb, Rb-Sr and K measurements on some Apollo 15 and Apollo 16 samples. *In **The Apollo 15 Lunar Samples*** (Chamberlain and Watkins, eds) 391-395. Lunar Planetary Institute, Houston.
- Taylor L.A., McCallister R.T. and Sardi O. (1973c) Cooling histories of lunar rocks based on opaque mineral geothermometers. *Proc. 4th Lunar Sci. Conf.* 819-828.

Taylor L.A., Uhlmann D.R., Hopper R.W. and Misra K.C. (1975b) Absolute cooling rates of lunar rocks: Theory and application. *Proc. 6th Lunar Sci. Conf.* 181-191.

Taylor L.A. and McCallister R.H. (1972) An experimental investigation of the significance of zirconium partitioning in lunar ilmenite and ulvospinel. *Earth Planet. Sci. Lett.* **17**, 105-109.

Unruh D.M., Stille P., Patchett P.J. and Tatsumoto M. (1984) Lu-Hf and Sm-Nd evolution in lunar mare basalts. *Proc. 14th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **88**, B459-B477.

Walker D., Longhi J., Lasaga A.C., Stolper E.M., Grove T.L. and Hays J.F. (1977) Slowly cooled microgabbros 15555 and 15065. *Proc. 8th Lunar Sci. Conf.* 1521-1547.

Wänke H., Palme H., Baddenhausen H., Dreibus G., Jagoutz E., Kruse H., Palme C., Spettel B., Teschke F. and Thacker R. (1975a) New data on the chemistry of lunar samples: Primary matter in the lunar highlands and the bulk composition of the moon. *Proc. 6th Lunar Sci. Conf.* 1313-1340.

Wark D.A., Reid A.F., Lovering J.F. and El Goresy A. (1973) Zirconolite (versus Zirkelite) in lunar rocks (abs). *Lunar Sci.* **IV**, 764-766.

Wolf R., Woodrow A. and Anders E. (1979) Lunar basalts and pristine highland rocks: Comparison of siderophile and volatile elements. *Proc. 10th Lunar Planet. Sci. Conf.* 2107-2130.

Yajima T. and Hafner S.S. (1974) Cation distribution and equilibrium temperature of pigeonite from basalt 15065. *Proc. 5th Lunar Sci. Conf.* 769-784.