

15556
Vesicular Mare Basalt
1542.3 grams

DRAFT



Figure 1: Photo of 15556,0. Largest vesicles are about 6 mm. NASA S87-48187.

Introduction

15556 was collected about 60 meters from the edge of Hadley Rille in an area called The Terrace. The lunar regolith was thin in this area, with abundant rock samples (basalts) exposed (Swann et al. 1971). The sample is ~3.4 b.y. old.

Petrography

15556 is a very vesicular, olivine-normative mare basalt. The vesicle size varies across the sample (4-8 mm), but the grain size of the minerals doesn't vary. McGee et al. (1977), Rhodes and Hubbard (1973) and

Ryder (1985) find that it is fine-grained with anhedral phenocrysts (0.4mm) set in a subophitic matrix of pyroxene, plagioclase and ilmenite tablets (0.4 to 0.8 mm). The mesostasis contains glass, silica, troilite and separate grains of iron.

The original sample catalog reports vesicles make up as much as 50% by volume of the sample (figures 1 and 8) and range from 1 to 5 mm in size. Garvin et al. (1972) and Goldberg et al. (1976) studied the vesicles, finding that they were even larger (average = 4.1 mm). Note similarity with sample 15529.




Figure 2: Photomicrograph of thin section of 15556 showing vesicle wall. Scale about 3 mm. NASA S71-52492. Note relatively high abundance of opaque phases.

Mineralogical Mode of 15556

	Sample Catalog Butler 1971	Rhodes and Hubbard 1973	McGee et al. 1979
Olivine	5	0.1	0.1
Pyroxene	50	57	57
Plagioclase	30	38	38
Ilmenite	3	2.1	2
Spinel	5	1	1
Mesostasis	1	1	1
Silica	5	0.8	1



Figure 3: Thin section photos of 15556 (same area as figure 2). a) plane polarized light, b) crossed polarizers. NASA S71-51762-3. Scale about 2 mm.

The original catalogue reported a large “xenocryst”.

Mineralogy

The mineralogy of 15556 has not been well studied. Some pyroxene analyses were reported by Brunfelt et al. (1972) and McGee et al. (1977), El Goresy et al. (1976) studied the chromite, von Englehardt (1979) ilmenite, Huffman et al. (1975) confirmed olivine using Mossbauer and Ryder (1985) found trace ‘ragged’ olivine and small clinopyroxene (figures 2 and 3). The independent iron grains apparently haven’t been analyzed.

Humphries et al. (1972) determined the experimental phase diagram showing that olivine was on the liquidus and arguing for olivine concentration from the original volcanic liquid.

Chemistry

Ryder and Shuraytz (2001) and others analyzed 15556 (figures 5 and 6). Moore et al. (1973) found only 13-16 ppm carbon in 15556. Gibson et al. (1975) reported

that the sulfur content was about the same as for 15555 – non vesicular equivalent.

Radiogenic age dating

Kirsten et al. (1972) determined a K/Ar age of 3.4 ± 0.1 b.y.

Cosmogenic isotopes and exposure ages

Rancitelli et al. (1972) determined the cosmic-ray-induced activity of $^{26}\text{Al} = 103$ dpm/kg, $^{22}\text{Na} = 40$ dpm/kg, $^{46}\text{Sc} = 6.5$ dpm/kg, $^{54}\text{Mn} = 41$ dpm/kg, and $^{56}\text{Co} = 11$ dpm/kg.

Other Studies

Kirsten et al. (1972) reported analyses for isotopic ratios of rare gases in 15556. Gibson and Andrawes (1978) studied the volatiles released by crushing the rock (nothing happened!). Nagata studied the magnetic properties.




Figure 4: Composition of pyroxene in 15556 (from McGee et al. 1977).

Processing

A slab was cut (figure 7). There are 29 thin sections (yet still no good petrographic description!).




Figure 5: Normalized rare-earth-element diagram for 15556 (data by Ryder and Shuraytz 2001). Compared with 15555.

Lunar Basalts




Figure 6: Composition of 15556 compared with other lunar basalts.

Table 1. Chemical composition of 15556.

reference weight	Rhodes 73	Wolf 79 Ganapathy73	Mason 72	Brunfeldt72	Strasheim72	Ryder2001	Rancitelli72						
SiO2 %	45.11 (a)		46.18 (c)		46.2 (f)	45.7 (a)	45.7 (a)						
TiO2	2.76 (a)		2.64 (c)	2.05 (e)	2.68 (f)	2.62 (a)	2.58 (a)						
Al2O3	9.43 (a)		9.85 (c)	9.24 (e)	9.44 (f)	9.48 (a)	9.24 (a)						
FeO	22.25 (a)		21.7 (c)	21.36 (e)	21.62 (f)	21.67 (a)	21.89 (a)	21.7 (e)	22.2 (e)				
MnO	0.29 (a)		0.32 (c)	0.3 (e)	0.26 (f)	0.28 (a)	0.28 (a)						
MgO	7.73 (a)		8.03 (c)		8.09 (f)	8.17 (a)	8.67 (a)						
CaO	10.83 (a)		10.72 (c)	9.93 (e)	10.54 (f)	10.56 (a)	10.38 (a)						
Na2O	0.26 (a)		0.3 (c)	0.28 (e)	0.22 (f)	0.257 (a)	0.273 (a)	0.27 (e)	0.27 (e)				
K2O	0.03 (a)		0.09 (c)		0.06 (f)	0.047 (a)	0.046 (a)				0.053 (h)		
P2O5	0.08 (a)		0.07 (c)		0.08 (f)	0.07 (a)							
S %	0.08 (a)												
sum													
Sc ppm				43.1 (e)	35 (f)			45.6 (e)	45.9 (e)				
V			165 (d)	266 (e)	255 (f)								
Cr			5200 (d)	3230 (e)	4310 (f)	5740 (a)	4452 (a)	5550 (e)	4430 (e)				
Co			46 (d)	50.3 (e)	49 (f)			50.2 (e)	48.9 (e)				
Ni			65 (d)	50 (e)	57 (f)	63 (a)	40 (a)	73 (e)	63 (e)				
Cu			10 (d)	7.1 (e)	9 (f)	6 (a)	7 (a)						
Zn	2.1 (b)			1.2 (e)									
Ga		5 (d)		3.7 (e)									
Ge ppb	9.8 (b)												
As				0.05 (e)									
Se	142 (b)			0.106 (e)									
Rb	0.1 (b)	<5 (d)		0.84 (e)	3 (f)	6 (a)	4 (a)						
Sr		102 (d)		88 (e)	96 (f)	101 (a)	99 (a)	126 (e)	111 (e)				
Y		50 (d)			32 (f)	24 (a)	24 (a)						
Zr		100 (d)			85 (f)	94 (a)	92 (a)						
Nb					8 (f)	10 (a)	9 (a)						
Mo													
Ru													
Rh													
Pd ppb													
Ag ppb	0.85 (b)		<7 (e)										
Cd ppb	28 (b)												
In ppb	0.56 (b)		<2 (e)										
Sn ppb													
Sb ppb	0.13 (b)												
Te ppb	2.7 (b)												
Cs ppm	0.03 (b)			0.032 (e)									
Ba		50 (d)		59 (e)	74 (g)			44 (e)	53 (e)				
La				4.8 (e)	3.7 (g)			5.2 (e)	5.32 (e)				
Ce				18 (e)	16 (g)			15.8 (e)	14.3 (e)				
Pr					2.6 (g)								
Nd					10.8 (g)			13 (e)	12 (e)				
Sm				4 (e)	3.95 (g)			3.65 (e)	3.85 (e)				
Eu				1 (e)	1.1 (g)			0.96 (e)	0.95 (e)				
Gd					4.5 (g)								
Tb				0.77 (e)	0.85 (g)			0.87 (e)	0.83 (e)				
Dy				4.4 (e)	5.3 (g)								
Ho				0.91 (e)	1.3 (g)								
Er				3.3 (e)	2.85 (g)								
Tm					0.48 (g)								
Yb				1.59 (e)	2.5 (g)			2.27 (e)	2.34 (e)				
Lu				0.39 (e)	0.33 (g)			0.31 (e)	0.33 (e)				
Hf				3.1 (e)				2.89 (e)	2.77 (e)				
Ta				0.4 (e)				0.39 (e)	0.43 (e)				
W ppb				430 (e)									
Re ppb	0.00413 (b)												
Os ppb													
Ir ppb	0.039 (b)		<0.1 (e)										
Pt ppb													
Au ppb	0.026 (b)			0.85 (e)									
Th ppm				0.4 (e)				0.46 (e)	0.42 (e)	0.56 (h)			
U ppm	0.145 (b)			0.21 (e)						0.15 (h)			

technique: (a) XRF, (b) RNAA, (c) wet chem., (d) OES, (e) INAA, (f) mixed, (g) MS, (h) radiation counting





Figure 7: Slab cut from 15556. Cube is 1 cm. NASA S81-37816.







Figure 8: Photo of 15556,188. NASA S97-16862. Large vesicles are about 8 mm.



References for 15556

Butler P. (1971) Lunar Sample Catalog, Apollo 15. Curators' Office, MSC 03209

Brunfelt A.O., Heier K.S., Nilssen B., Steiennes E. and Sundvoll B. (1972) Elemental composition of Apollo 15 samples. In **The Apollo 15 Lunar Samples** (Chamberlain J.W. and Watkins C., eds.), 195-197. Lunar Science Institute, Houston.

El Goresy A., Prinz M. and Ramdohr P. (1976a) Zoning in spinels as an indicator of the crystallization histories of mare basalts. *Proc. 7th Lunar Sci. Conf.* 1261-1279.

Engelhardt W. von (1979) Ilmenite in the crystallization sequence of lunar rocks. *Proc. 10th Lunar Planet. Sci. Conf.* 677-694.

Humphries D.J., Biggar G.M and O'Hara M.J. (1972) Phase equilibria and origin of Apollo 15 basalts etc. In **The Apollo 15 Lunar Samples**. 103-107. Lunar Planetary Institute, Houston.

Kirsten T., Deubner J., Horn P., Kaneoka I., Kiko J., Schaeffer O.A. and Thio S.K. (1972) The rare gas record of Apollo 14 and 15 samples. *Proc. 3rd Lunar Sci. Conf.* 1865-1889.

Ganapathy R., Morgan J.W., Krahenbuhl U. and Anders E. (1973) Ancient meteoritic components in lunar highland rocks: Clues from trace elements in Apollo 15 and 16 samples. *Proc. 4th Lunar Sci. Conf.* 1239-1261.

Garvin J.B., Head J.W. and Wilson L. (1982) Magma vesiculation in Apollo 15 mare basalts: Observations and theory (abs). *Lunar Planet. Sci.* XIII 255-256.

Gibson E.K., Chang S., Lennon K., Moore G.W. and Pearce G.W. (1975a) Sulfur abundances and distributions in mare basalts and their source magmas. *Proc. 6th Lunar Sci. Conf.* 1287-1301.

Gibson E.K. and Andrawes F.F. (1978a) Nature of the gases released from lunar rocks and soils upon crushing. *Proc. 9th Lunar Planet. Sci. Conf.* 2433-2450.

Goldberg R.H., Trombrello T.A. and Burnett D.S. (1976a) Fluorine as a constituent in lunar magmatic gases. *Proc. 7th Lunar Sci. Conf.* 1597-1613.

LSPET (1972a) The Apollo 15 lunar samples: A preliminary description. *Science* **175**, 363-375.

Mason B., Jarosewich E., Melson W.G. and Thompson G. (1972) Mineralogy, petrology, and chemical composition of lunar samples 15085, 15256, 15271, 15471, 15475,

15476, 15535, 15555, and 15556. *Proc. 3rd Lunar Sci. Conf.* 785-796.

McGee P.E., Warner J.L. and Simonds C.H. (1977) Introduction to the Apollo Collections. Part I: Lunar Igneous Rocks. Curators Office, JSC.

Moore C.B., Lewis C.F., and Gibson E.K. (1972) Carbon and nitrogen in Apollo 15 lunar samples. In *The Apollo 15 Lunar Samples* (Chamberlain J.W. and Watkins C., eds.), 316-318. The Lunar Science Institute, Houston.

Moore C.B., Lewis C.F. and Gibson E.K. (1973) Total carbon contents of Apollo 15 and 16 lunar samples. *Proc. 4th Lunar Sci. Conf.* 1613-1923.

Rancitelli L.A., Perkins R.W., Felix W.D. and Wogman N.A. (1972) Lunar surface processes and cosmic ray characterization from Apollo 12-15 lunar samples analyses. *Proc. 3rd Lunar Sci. Conf.* 1681-1691.

Rhodes J.M. and Hubbard N.J. (1973) Chemistry, Classification, and petrogenesis of Apollo 15 mare basalts. *Proc. 4th Lunar Sci. Conf.* 1127-1148.

Ryder G. (1985) Catalog of Apollo 15 Rocks (three volumes). Curatorial Branch Pub. # 72, JSC#20787

Ryder G. and Schuraytz B.C. (2001) Chemical variations of the large Apollo 15 olivine-normative mare basalt rock samples. *J. Geophys. Res.* **106**, E1, 1435-1451.

Schwerer F.C. and Nagata T. (1976) Ferromagnetic-superparamagnetic granulometry of lunar surface materials. *Proc. 7th Lunar Sci. Conf.* 759-778.

Sutton R.L., Hait M.H., Larson K.B., Swann G.A., Reed V.S. and Schaber G.G. (1972) Documentation of Apollo 15 samples. Interagency report: *Astrogeology* 47. USGS

Swann G.A., Hait M.H., Schaber G.C., Freeman V.L., Ulrich G.E., Wolfe E.W., Reed V.S. and Sutton R.L. (1971) Preliminary description of Apollo 15 sample environments. U.S.G.S. Interagency report: 36. pp219 with maps

Swann G.A., Bailey N.G., Batson R.M., Freeman V.L., Hait M.H., Head J.W., Holt H.E., Howard K.A., Irwin J.B., Larson K.B., Muehlberger W.R., Reed V.S., Rennilson J.J., Schaber G.G., Scott D.R., Silver L.T., Sutton R.L., Ulrich G.E., Wilshire H.G. and Wolfe E.W. (1972) 5. Preliminary Geologic Investigation of the Apollo 15 landing site. In Apollo 15 Preliminary Science Rpt. NASA SP-289. pages 5-1-112.

Wolf R., Woodrow A. and Anders E. (1979) Lunar basalts and pristine highland rocks: Comparison of siderophile and

volatile elements. *Proc. 10th Lunar Planet. Sci. Conf.* 2107-2130.