

60025
Ferroan Anorthosite
 1836 grams



Figure 1: 60025. NASA #S72-41586. Cube and scale are 1 cm. Note the thick black glass coating and numerous micrometeorite pits.

Introduction

Lunar sample 60025 is a large sample of coarse-grained cataclastic anorthosite (figure 1). Or, rather, it is a mixture of pieces from a related sequence of anorthosites, because the mafic minerals vary in composition (Ryder 1982, James et al. 1991). Warren and Wasson (1977) termed 60025 “pristine” because of its coarse grain size and low meteoritic siderophiles (Ni, Ir). The age of 60025 has been determined by Pb-

Pb and Sm-Nd to be 4.36 b.y. and it has very low $^{87}\text{Sr}/^{86}\text{Sr}$ (0.699).

Ferroan anorthosites are one of the major lunar rock types (Dowty et al. 1974, James 1980) and are thought to be pieces of the original lunar crust formed by plagioclase floatation from a moon-wide magma ocean (e.g. Smith et al. 1970, Wood et al. 1970, Warren 1990).

Mineralogical Mode of 60025

	Dixon and Papike 1975	Warren and Wasson 1978	James et al. 1991	James et al. mafic region
Plagioclase	98 vol. %	70	74	5
Olivine		20	20	45
Orthopyroxene		10	5	45
Augite			1	5
Chromite			0.05	0.2
Ilmenite			0.02	



Figure 2: Group photo of pieces cut from first slab. Note the chalky white interior of 60025 and the patina on the exterior surface. Cubes are about 1 cm. NASA # S72-49109.

Alternatively, 60025 may be from a layered igneous pluton emplaced within the lunar crust (James 1980).

Ryder and Norman (1980) provided a compilation of what was known about 60025 up to that time. James et al. (1991) studied the mafic portions of 60025.

60025 was collected near the Lunar Module and its orientation was documented by photography.

Petrography

60025 has been described as a moderately shocked, cataclastic anorthosite (Hodges and Kushiro 1973, Walker et al. 1973, Dixon and Papike 1975, **Ryder 1982**, James et al. 1991) made up of mostly calcic plagioclase (see mode). The first slabs cut from 60025 were nearly pure plagioclase (figure 2), but other portions of 60025 were found to be more mafic (see Warren and Wasson 1977, Ryder 1982, James et al. 1991). Most pyroxene grains are distinct augite or orthopyroxene, but some grains are exsolved into end member components (figure 3). There are trace amounts of silica, ilmenite and chromite.

A variety of textures are reported (figure 3). In some areas, large fractured plagioclase grains (2-4 mm) sit in a sea of crushed plagioclase and pyroxene (Walker et al. 1973).

Pyroxene compositions are variable in different portions of the rock (En_{50} to En_{70}) indicating a differentiation sequence (Ryder 1982). However, the origin of some pyroxene by sub-solidus exsolution should be considered (see Smith and Steele 1974). The wide composition gap between augite and hypersthene (Wo_{45} to Wo_2) indicates thermal equilibrium well below magmatic temperatures (Walker et al 1973, Dixon and Papike 1975), presumably as a slowly-cooled plutonic rock.

Mineralogy

Plagioclase: A number of people have analyzed the plagioclase in 60025 and found it to be uniform and very calcic (An_{96-98}) (see table for some of the analyses).




Figure 3: Plane-polarized and cross-polarized photos of thin section 60025,21. Field of view is 2.5 mm. S79-27300 and 301.

Olivine: Warren and Wasson (1977), Ryder (1982) and James et al. (1991) find that olivine in various portions of 60025 has a range of compositions (Fo₄₂₋₆₆).

Pyroxene: Pyroxene is present as subhedral to anhedral grains included in plagioclase, as anhedral grains at grain boundaries and in mafic clumps with olivine. 60025 contains both high-Ca and low-Ca pyroxene with a wide miscibility gap (figure 4). Hodges and Kushiro (1973) noted that three pyroxenes (augite, hypersthene and pigeonite) were probably present at time of crystallization. Pigeonite is now exsolved to endmember compositions. Fine exsolution is also found in sub-calcic augite (Hodges and Kushiro 1973). Dixon and Papike (1975) find that the composition of coexisting augite and orthopyroxene indicates an equilibrium temperature of less than 800 deg. C (see also Lindsley et al. 1974). Floss et al. (1998) determined the REE for pyroxenes in 60025 (figure 6).

The variation of FeO/MgO in pyroxene, from one portion of 60025 to another, is the evidence used by Ryder (1982) to argue that 60025 may actually be a mix of anorthosites from a differentiated sequence.

Opagues: Hodges and Kushiro (1973) and James et al. (1991) determined the composition of chromite and ilmenite.

Glass: The black glass found attached to the surface of 60025 has been studied by See et al. (1986) and Morris et al. (1986).

Chemistry

Plagioclase dominates the composition of 60025 (figures 6 and 9). Haskin et al. (1973, 1981) and Krahenbuhl et al. (1973) showed that 60025 was free of meteoritic siderophile elements (Ni, Ir). James et al. (1991) determined the composition of mafic portions of 60025 (table 1).

Radiogenic age dating

Tera and Wasserburg (1972), Papanastassiou and Wasserburg (1972), Nunes et al. (1974), Schaeffer and Husain (1974), Hanan and Tilton (1987) and Carlson and Lugmair (1988) have attempted to date 60025. The ⁸⁷Sr/⁸⁶Sr ratio is extremely (0.699) low indicating that 60025 is very old and formed from a low Rb source (Nyquist et al. 1974). The mafic minerals allowed a precise Sm/Nd isochron (figure 7), which is only about 110 m.y. younger than the age of the moon. However, the interpretation that 60025 is in fact a mix of related anorthosites, this is perhaps equivalent to a “whole rock” isochron.

The exact age of 60025 has been recently (2011) reinvestigated by Pb/Pb and Nd/Sm and found to be significantly younger (4.36 b.y.) than previously thought (Borg et al. 2011).

Cosmogenic isotopes and exposure ages

Kurt Marti determined the exposure age of 60025 as 1.9 m.y. (see Drozd et al. 1977 and Arvidson et al. 1975). Fruchter et al. (1977) and Kohl et al. (1977) determined ⁵³Mn in 60025 as 105 ± 13 and 76 ± 6 dpm/kg respectively.






Figure 4: Pyroxene data from various investigators studying various different regions of 60025.



Other Studies

Sato (1986) determined the intrinsic oxygen fugacity of 60025. Flory et al. (1973) studied the hydrocarbons that were emitted during heating of 60025 (apparently produced by hydrolysis of solar wind implanted carbon). Schaeffer and Husain (1974), Lightner and Marti (1974) and Leich and Niemeyer (1975) analyzed the rare gases in 60025. Katsube and Collett (1973), Gold et al. (1976) and Cisowski et al. (1976) determined electrical and magnetic properties. Hapke et al. (1978) collected ultraviolet reflectance spectra. Simmons et al. (1975) studied the microcracks and Jeanloz and Ahrens (1978) studied the compressibility (equation of state). Sondergeld et al. (1979) determined seismic wave velocities to compare with seismic data collected at Apollo 16.

Processing

Two thick slabs have been cut from 60025 (figures 2 and 10) and several pieces are used in public displays (see flow diagram). Mafic-rich pieces were taken from the side opposite the slabs (W1).

List of Photo #s	
S72-41586-41591	color mug shots
S72-42582-42597	B&W exterior
S72-49096-49109	first slab
S72-49095	slab assembled
S72-49094	end piece, subdivided
S74-28116	second slab




Figure 5: Plagioclase-low-Ca pyroxene diagram for various portions of ferroan anorthosite 60025.

Plagioclase in 60025

electron probe (wt.%)

	Papike 97	Hanson 79	McGee 93	Dixon and Papike 1975				James et al. 1991		
SiO ₂	43.9			43.4	44.7	44.8	44.4	44.8	44.15	44.08
Al ₂ O ₃	35.3			35.2	35.2	35.2	35.4	34.4	36.03	36.01
FeO	0.18	0.145	0.145	0.19	0.15	0.19	0.17	0.23		
MgO	0.051	0.064	0.077	0.04	0.02	0.06	0.07	0.02	0.058	0.048
CaO	19			19.5	19.6	19.3	19.5	19.6	19.13	19.08
Na ₂ O	0.426			0.42	0.39	0.48	0.46	0.36	0.34	0.359
K ₂ O	0.009	0.031		0	0	0	0	0	0.01	0.06
Ab	3.9	3.9								
An	96		96.9	96.3	96.5	95.7	95.9	96.8	96.82	96.65
Or	0.057									

ion probe (ppm)

	Meyer 79	Steele 80	Papike 97	Floss 98	
Li	1.4	1.9			
Mg	474	410			mafic area
Ti	85	80			
Sr	197	240	168		
Y			0.22		
Ba	15	15	10.42		
La			0.29	0.24	0.33
Ce			0.483	0.67	0.82
Nd			0.26	0.39	0.43
Sm			0.086	0.094	0.06
Eu			1.16	1.03	1.05
Gd			0.066	0.093	0.071
Dy			0.047	0.055	0.039
Er			0.018	0.025	0.018
Yb			0.016	0.031	0.022




Figure 6: Normalized rare-earth-element diagram for 60025 and its minerals. Data for whole rock from Haskin et al. 1981, data for minerals from Floss et al. 1998. Note that plagioclase dominates.




Figure 7: Sm-Nd mineral isochron for 60025 by Carlson and Lugmair (1988).




Figure 8a: Argon release pattern for 60025 by Schaeffer and Husain (1975).




Figure 8b: Sm-Nd mineral isochrons for portion of 60025 (Borg et al. 2011).




Figure 8c: Pb-Pb isochron for 60025 (Borg et al. 2011).

Summary of Age Data for 60025

	Pb/Pb	Sm/Nd	Ar/Ar
Schaeffer and Husain 1974			4.21 ± 0.06 b.y. 4.17 ± 0.06
Hanan and Tilton 1987			
Plagioclase	4.51 ± 0.01 b.y.		
Mafic	4.42		
Carlson and Lugmair 1988		4.44 ± 0.02	
Borg et al. 2011	4.359 ± 0.002	4.367 ± 0.011	

Table 1a. Chemical composition of 60025.

reference weight	Wiesmann 75						Ehmann 75	
	Wiesmann 75	Rose 73	Haskin 73	Laul 73	Krahenbuhl 73 Wolf 79	Nakamura 73	Garg 76	Janghorbani 73
SiO ₂ %		43.03	(b) 45.3	(c)		44.75		42.79
TiO ₂	0.02	(a) 0.02	(b) <0.02	(c) <0.06	(c)	0.2		
Al ₂ O ₃		35.21	(b) 34.2	(c) 35.5	(c)	35.28		36.84
FeO		0.67	(b) 0.495	(c) 0.35	(c)	0.31		1.3
MnO		0.03	(b) 0.008	(c) 0.009	(c)	0.013		0.012
MgO	0.14	(a) 0.27	(b) 0.206	(c)		0.17		0.16
CaO		18.92	(b) 19.8	(c) 18.2	(c)	19.38		17.2
Na ₂ O		0.49	(b) 0.45	(c) 0.414	(c)	0.44		0.44
K ₂ O	0.008	(a) 0.03	(b) 0.113	(c) <0.01	(c)	0.028		
P ₂ O ₅		0				0.003		
S %								
sum								
Sc ppm			0.7	(c) 0.55	(c)			0.4
V		<5	(b)	<7	(c)			
Cr		0	(b) 24	(c) 102	(c)	205		
Co		3.6	(b) 0.73	(c) 0.7	(c)			0.7
Ni		30	(b) 1.1	(c)		<3	(d)	
Cu		8.4	(b)					
Zn		15	(b) <2	(c)		0.17	(d)	
Ga			4	(c)				
Ge ppb						2.3	(d)	
As								
Se						0.022	(d)	
Rb	0.03	(a) 0.8	(b) <0.1	(c)		0.017	(d)	
Sr	231	(a) 205	(b)					
Y								
Zr	9	(a) <10	(b)					0.5
Nb		<10	(b)					
Mo								
Ru								
Rh								
Pd ppb								
Ag ppb						0.22	(d)	
Cd ppb						7.25	(d)	
In ppb								
Sn ppb								
Sb ppb						0.035	(d)	
Te ppb						65	(d)	
Cs ppm			<0.003	(c)		0.002	(d)	
Ba	13.2	(a) 27	(b)	10	(c)	15.87	(a)	
La	0.294	(a)	0.28	(c) 0.3	(c)	0.301	(a)	
Ce	0.607	(a)	0.65	(c) 0.8	(c)	0.645	(a)	
Pr								
Nd	0.368	(a)	0.42	(c) 0.5	(c)	0.361	(a)	
Sm	0.091	(a)	0.092	(c) 0.12	(c)	0.0881	(a)	
Eu	1.13	(a)	1.04	(c) 1.04	(c)	1.044	(a) 1.1	
Gd	0.103	(a)				0.0895	(a)	
Tb			0.2	(c) <0.02	(c)			
Dy	0.083	(a)	0.19	(c) 0.11	(c)	0.0783	(a)	
Ho								
Er	0.042	(a)	0.05	(c)		0.042	(a)	
Tm								
Yb	0.042	(a)	0.048	(c) 0.051	(c)	0.0429	(a)	
Lu	0.0046	(a)	0.006	(c) 0.005	(c)	0.00465	(a)	
Hf			0.02	(c) 0.013	(c)			0.0165
Ta				<0.01	(c)			
W ppb								
Re ppb								
Os ppb								
Ir ppb								
Pt ppb								
Au ppb								
Th ppm								
U ppm						0.00092	(d)	

technique (a) IDMS, (b) combined XRF, microchem., emiss. spec., (c) INAA, (d) RNAA

Table 1b. Chemical composition of 60025.

reference weight	glass						Haskin 82 av. 12	mafic patches				
	Morris 86	Haskin 81						James 91 ,702WR ,691WR ,697WR				
SiO ₂ %	44.54	(b)										
TiO ₂	0.36	(b)										
Al ₂ O ₃	27.65	(b)										
FeO	5.14	(b)	0.32	0.362	0.36	1.31	0.315	0.63	(a)	0.665	13.1	11.8
MnO												
MgO	6.35	(b)										
CaO	15.46	(b)	18.5	18.2	18.1	17.8	18.4	18.6	(a)	18.5	8.3	10.4
Na ₂ O	0.34	(b)	0.45	0.42	0.43	0.4	0.44	0.42	(a)	0.423	0.169	0.219
K ₂ O	0.08	(b)										
P ₂ O ₅												
S %												
sum												
Sc ppm	6.15	(a)	0.47	0.47	0.55	2.47	0.54	1.02	(a)	1.05	9.66	9.18
V												
Cr	754	(a)	0.55	0.69	0.59	1.99	0.54	73	(a)	42.7	812	781
Co	49	(a)	14.7	13.4	14.7	8	12.9	1.05	(a)	1.07	23.8	21.2
Ni	711	(a)										
Cu												
Zn												
Ga												
Ge ppb												
As												
Se												
Rb												
Sr			226	208	207	193	198			199	71	39
Y												
Zr												
Nb												
Mo												
Ru												
Rh												
Pd ppb												
Ag ppb												
Cd ppb												
In ppb												
Sn ppb												
Sb ppb												
Te ppb												
Cs ppm												
Ba	233	(a)	10.4	12	11.5	10	65			16		
La	10.49		0.289	0.252	0.234	0.23	0.275	0.27		0.235	0.14	0.173
Ce	31.3		0.67	0.69	0.54		0.7	0.65		0.62	0.71	
Pr												
Nd												
Sm	4.79		0.097	0.088	0.085	0.089	0.093	0.098		0.086	0.089	0.086
Eu	1.13		1.11	1.05	1.07	1	1.01	1.025		0.97	0.46	0.586
Gd												
Tb	0.97		0.0183	0.0152	0.0178	0.0252	0.015			0.018	0.029	0.015
Dy												
Ho												
Er												
Tm												
Yb			0.04	0.045	0.049	0.085	0.04	0.051		0.032	0.211	0.178
Lu	0.45		0.0046	0.0052	0.005	0.0111		0.0081		0.009	0.037	0.031
Hf	3.53		0.0078	0.0064	0.0086	0.021	0.018				0.076	0.08
Ta	0.43										0.018	0.036
W ppb												
Re ppb												
Os ppb												
Ir ppb												
Pt ppb												
Au ppb												
Th ppm	2.71											
U ppm	0.22											
technique			(a) INAA, (b) microprobe									




Figure 9: Chemical composition of black glass coating on 60025, compared with composition of interior anorthosite.




Figure 10: Second slab cut from 60025 in 1974. Scale is in cm and cube is 1 inch. Slab was about 1 inch thick. NASA photo # S74-28116





Figure 11: Position of saw blade prior to cutting first slab. White line shows approximate position of second slab. Cube is 1 inch. S76-21590 (copy of processing photo).







Figure 12: Processing photo of second slab. Big cube is 1 inch. S74-28115. Note the saw marks.



References for 60025

- Arvidson R., Crozaz G., Drozd R.J., Hohenberg C.M. and Morgan C.J. (1975) Cosmic ray exposure ages of features and events at the Apollo landing sites. *The Moon* **13**, 259-276.
- Dixon J.R. and Papike J.J. (1975) Petrology of anorthosites from the Descartes region of the moon: Apollo 16. *Proc. 6th Lunar Sci. Conf.* 263-291.
- Dowty E., Prinz M. and Keil K. (1974b) Ferroan anorthosite: a widespread and distinctive lunar rock type. *Earth Planet. Sci. Lett.* **24**, 15-25.
- Drozd R.J., Hohenberg C.M., Morgan C.J. and Ralston C.E. (1974) Cosmic-ray exposure history at the Apollo 16 and other lunar sites: lunar surface dynamics. *Geochim. Cosmochim. Acta* **38**, 1625-1642.
- Borg L.E., Connelly J.N., Boyet M. and Carlson R.W. (2011) Chronological evidence that the Moon is either young or did not have a global magma ocean. *Nature* **477**, 70 – 72.
- Carlson R.W. and Lugmair G.W. (1988) The age of ferroan anorthosite 60025: oldest crust on a young Moon? *Earth Planet. Sci. Lett.* **90**, 119-130.
- Ehmann W.D. and Chyi L.L. (1974) Abundances of the group IVB elements, Ti, Zr, and Hf and implications of their ratios in lunar materials. *Proc. 5th Lunar Sci. Conf.* 1015-1024.
- Ehmann W.D., Chyi L.L., Garg A.N., Hawke B.R., Ma M.-S., Miller M.D., James W.D. and Pacer R.A. (1975a) Chemical studies of the lunar regolith with emphasis on zirconium and hafnium. *Proc. 6th Lunar Sci. Conf.* 1351-1361.
- Floss C., James O.B., McGee J.J. and Crozaz G. (1998) Lunar ferroan anorthosite petrogenesis: Clues from trace distributions in FAN subgroups. *Geochim. Cosmochim. Acta* **62**, 1255-1283.
- Fruchter J.S., Rancitelli L.A., Laul J.C. and Perkins R.W. (1977) Lunar regolith dynamics based on analysis of the cosmogenic radionuclides ²²Na, ²⁶Al and ⁵³Mn. *Proc. 8th Lunar Planet. Sci. Conf.* 3595-3605.
- Garg A.N. and Ehmann W.N. (1976a) Zr-Hf fractionation in chemically defined lunar rock groups. *Proc. 7th Lunar Sci. Conf.* 3397-3410.
- Gold T., Bilson E. and Baron R.L. (1976a) The surface chemical composition of lunar samples and its significance for optical properties. *Proc. 7th Lunar Sci. Conf.* 901-911.
- Hanan B.B. and Tilton G.R. (1987) 60025: relict of primitive lunar crust? *Earth Planet. Sci. Lett.* **84**, 15-21.
- Hapke B.W., Partlow W.D., Wagner J.K. and Cohen A.J. (1978) Reflectance measurements of lunar materials in the vacuum ultraviolet. *Proc. 9th Lunar Planet. Sci. Conf.* 2935-2947.
- Hansen E.C., Steele I.M. and Smith J.V. (1979a) Lunar highland rocks: Element partitioning among minerals 1: Electron microprobe analyses of Na, K, and Fe in plagioclase; mg partitioning with orthopyroxene. *Proc. 10th Lunar Planet. Sci. Conf.* 627-638.
- Haskin L.A., Helmke P.A., Blanchard D.P., Jacobs J.W. and Telunder K. (1973) Major and trace element abundances in samples from the lunar highlands. *Proc. 4th Lunar Sci. Conf.* 1275-1296.
- Haskin L.A., Lindstrom M.M., Salpas P.A. and Lindstrom D.L. (1981) On compositional variations among lunar anorthosites. *Proc. 12th Lunar Planet. Sci. Conf.* 41-66.
- Hodges F.N. and Kushiro I. (1973) Petrology of Apollo 16 lunar highland rocks. *Proc. 4th Lunar Sci. Conf.* 1033-1048.
- James O.B., Lindstrom M.M. and McGee J.J. (1991) Lunar ferroan anorthosite 60025: Petrology and chemistry of mafic lithologies. *Proc. 21st Lunar Planet. Sci. Conf.* 63-87. Lunar Planetary Institute, Houston.
- James O.B. (1980) Rocks of the early lunar crust. *Proc. 11th Lunar Planet. Sci. Conf.* 365-393.
- Janghorbani M., Miller M.D., Ma M.-S., Chyi L.L. and Ehmann W.D. (1973) Oxygen and other elemental abundance data for Apollo 14, 15, 16 and 17 samples. *Proc. 4th Lunar Sci. Conf.* 1115-1126.
- Jeanloz R.F. and Ahrens T.J. (1978) The equation of state of a lunar anorthosite: 60025. *Proc. 9th Lunar Sci. Conf.* 2789-2803.
- Katsube T.J. and Collett L.S. (1973a) Electrical characteristics of Apollo 16 lunar samples. *Proc. 4th Lunar Sci. Conf.* 3101-3110.
- Kohl C.P., Murell M.T., Russ G.P. III and Arnold J.R. (1978) Evidence for the constancy of the solar cosmic ray flux over the past ten million years: ⁵³Mn and ²⁶Al measurements. *Proc. 9th Lunar Planet. Sci. Conf.* 2299-2310.
- Krahenbuhl U., Ganapathy R., Morgan J.W. and Anders E. (1973b) Volatile elements in Apollo 16 samples: Implications for highland volcanism and accretion history of the moon. *Proc. 4th Lunar Sci. Conf.* 1325-1348.

- Laul J.C. and Schmitt R.A. (1973) Chemical composition of Apollo 15, 16, and 17 samples. *Proc. 4th Lunar Sci. Conf.* 1349-1367.
- Lindsley D.H., King H.E. and Turnock A.C. (1974) Composition of synthetic agugite and hypersthene coexisting at 810 deg C: Application to pyroxenes from the lunar highland rocks. *Geophys. Res. Lett.* **1**, 134-136.
- Lightner B.D. and Marti K. (1974) Lunar trapped xenon. *Proc. 5th Lunar Sci. Conf.* 2023-2031.
- Leich D.A. and Niemeyer S. (1975) Trapped xenon in lunar anorthosite breccia 60015. *Proc. 6th Lunar Sci. Conf.* 1953-1965.
- McGee J.J. (1993) Lunar ferroan anorthosites: Mineralogy, compositional variations and petrogenesis. *J. Geophys. Res.* **98**, 9089-9105.
- Meyer C. (1979) Trace elements in plagioclase from the lunar highlands. In Papers presented to the **Conference on the Lunar Highlands Crust** (abs). LPI Contr. 394, 111-113. Lunar Planetary Institute, Houston
- Miller M.D., Pacer R.A., Ma M.-S., Hawke B.R., Lookhart G.L. and Ehmann W.D. (1974) Compositional studies of the lunar regolith at the Apollo 17 site. *Proc. 5th Lunar Sci. Conf.* 1079-1086.
- Moore C.B., Lewis C.F. and Gibson E.K. (1973) Total carbon contents of Apollo 15 and 16 lunar samples. *Proc. 4th Lunar Sci. Conf.* 1613-1923.
- Morris R.V., See T.H. and Horz F. (1986) Composition of the Cayley Formation at Apollo 16 as inferred from impact melt splashes. *Proc. 17th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **90**, E21-E42.
- Nakamura N., Masuda A., Tanaka T. and Kurasawa H. (1973) Chemical compositions and rare-earth features of four Apollo 16 samples. *Proc. 4th Lunar Sci. Conf.* 1407-1414.
- Nunes P.D., Tatsumoto M., Knight R.J., Unruh D.M. and Doe B.R. (1973b) U-Th-Pb systematics of some Apollo 16 lunar samples. *Proc. 4th Lunar Sci. Conf.* 1797-1822.
- Nyquist L.E., Bansal B.M. and Wiesmann H. (1975a) Rb-Sr ages and initial 87Sr/86Sr for Apollo 17 basalts and KREEP basalt 15386. *Proc. 6th Lunar Sci. Conf.* 1445-1465.
- Papanastassiou D.A. and Wasserburg G.J. (1972b) Rb-Sr age of a crystalline rock from Apollo 16. *Earth Planet. Sci. Lett.* **16**, 289-298.
- Papike J.J., Fowler G.W. and Shearer C.K. (1997) Evolution of the lunar crust: SIMS study of plagioclase from ferroan anorthosites. *Geochim. Cosmochim. Acta* **61**, 2343-2350.
- Rose H.J., Cuttitta F., Berman S., Carron M.K., Christian R.P., Dwornik E.J., Greenland L.P. and Ligon D.T. (1973) Compositional data for twenty-two Apollo 16 samples. *Proc. 4th Lunar Sci. Conf.* 1149-1158.
- Ryder G. and Norman M.D. (1979b) Catalog of pristine non-mare materials Part 2. Anorthosites. Revised. Curators Office JSC #14603
- Ryder G. and Norman M.D. (1980) Catalog of Apollo 16 rocks (3 vol.). Curator's Office pub. #52, JSC #16904
- Ryder G. (1982b) Lunar anorthosite 60025, the petrogenesis of lunar anorthosites and the composition of the Moon. *Geochim. Cosmochim. Acta* **46**, 1591-1601.**
- Taylor H.P. and Epstein S. (1973) O¹⁸/O¹⁶ and Si³⁰/Si²⁸ studies of some Apollo 15, 16 and 17 samples. *Proc. 4th Lunar Sci. Conf.* 1657-1679.
- Sato M. (1976b) Oxygen fugacity values of some Apollo 16 and 17 rocks (abs). *Lunar Sci.* **VII**, 758-760. Lunar Planetary Institute, Houston.
- See T.H., Horz F. and Morris R.V. (1986) Apollo 16 impact-melt splashes: Petrography and major-element composition. *Proc. 17th Lunar Planet. Sci. Conf.* in *J. Geophys. Res.* **91**, E3-E20.
- Schaeffer O.A. and Husain L. (1974a) Chronology of lunar basin formation and ages of lunar anorthositic rocks (abs). *Lunar Sci.* **V**, 663-665.
- Schaeffer O.A. and Husain L. (1974b) Chronology of lunar basin formation. *Proc. 5th Lunar Sci. Conf.* 1541-1555.
- Simmons G., Siegfried R. and Richter D. (1975a) Characteristics of microcracks in lunar samples. *Proc. 6th Lunar Sci. Conf.* 3227-3254.
- Smith J.V., Anderson A.T., Newton R.C., Olsen E.J. and Wyllie P.J. (1970) A petrologic model for the moon based on petrogenesis, experimental petrology and physical properties. *J. Geol.* **78**, 381-405.
- Steele I.M., Hutcheon I.D. and Smith J.V. (1980) Ion microprobe analysis and petrogenetic interpretations of Li, Mg, Ti, K, Sr, Ba in lunar plagioclase. *Proc. 11th Lunar Planet. Sci. Conf.* 571-590.
- Tera F., Papanastassiou D.A. and Wasserburg G.J. (1974a) Isotopic evidence for a terminal lunar cataclysm. *Earth Planet. Sci. Lett.* **22**, 1-21.

Walker D., Longhi J., Grove T., Stolper E. and Hays J.F. (1973) Experimental petrology and origin of rocks from the Descartes Highlands. *Proc. 4th Lunar Sci. Conf.* 1013-1032.

Warren P.H. (1990) Lunar anorthosites and the magma ocean hypothesis: Importance of FeO enrichment in the parent magma. *Am. Mineral.* **75**, 46-58.

Warren P.H. and Wasson J.T. (1977) Pristine nonmare rocks and the nature of the lunar crust. *Proc. 8th Lunar Sci. Conf.* 2215-2235.

Warren P.H. and Wasson J.T. (1978) Compositional-petrographic investigation of pristine nonmare rocks. *Proc. 9th Lunar Planet. Sci. Conf.* 185-217.

Wasson J.T., Warren P.H., Kallemeyn G.W., McEwing C.E., Mitdefehldt D.W. and Boynton W.V. (1977) SCCRV, a major component of highlands rocks. *Proc. 8th Lunar Sci. Conf.* 2237-2252.

Wiesmann H. and Hubbard N.J. (1975) A compilation of the Lunar Sample Data Generated by the Gast, Nyquist and Hubbard Lunar Sample PI-Ships. Unpublished. JSC

Wolf R., Woodrow A. and Anders E. (1979) Lunar basalts and pristine highland rocks: Comparison of siderophile and volatile elements. *Proc. 10th Lunar Planet. Sci. Conf.* 2107-2130.

Wood J.A., Dickey J.S., Marvin U.B. and Powell B.N. (1970a) Lunar anorthosites. *Science* **167**, 602-604.

Wood J.A., Dickey J.S., Marvin U.B. and Powell B.N. (1970b) Lunar anorthosites and a geophysical model of the Moon. *Proc. Apollo 11 Lunar Sci. Conf.* 965-988.