

72275

Fragmental Polymict Breccia

3640 grams



Figure 1: Photo of pieces of 72275. Note thin patina and surface exposure of “Marble Cake” clast. NASA# S73-16077. Small cube is 1 cm.

Introduction

Lunar sample 72275 is a friable feldspathic breccia with an aphanitic matrix and several important clasts (figure 1). It was collected from the top of a layered boulder (#1), located within a landslide from the South Massif at the Apollo 17 site (Marvin 1975, Schmitt 1975) and is generally thought to represent ejecta from the Serenitatis basin (Dalrymple and Ryder 1996) (however, see evidence to the contrary in the paper by Morgan et al. 1975). The boulder may have rolled down the slope of the South Massif after the emplacement of the landslide, but there are no boulder tracks visible today so it is difficult to tell where exactly it came from (Wolfe 1975).

72275 was specifically collected from boulder 1, because it appeared to be representative of the matrix of the boulder. It was found that the chemical composition of 72275 matrix had higher trace element content than for the other three samples of this boulder (figure 2). This sample has a wide variety of clast types

derived from the lunar highlands (Stoeser *et al.* 1974, Ryder et al. 1977, Salpas et al. 1987, 1988). It has a high trace element content due to a high abundance of KREEPy non-mare basalt. It doesn’t contain any of the high-Ti mare material from the valley floor and is




Figure 2: Composition of matrix of 72275 with that of other samples from same boulder (#1). Data from Blanchard et al. (1975).




Figure 3: Photomicrographs of same area of thin section 72275,148: a) transmitted light, b) polarized light, c) reflected light. Note basalt clast in corner. Note also the porosity of matrix; best seen in reflected light. Scale is 1.4 mm across.

not a regolith breccia. The matrix and many of the clasts contain significant Ir and Au contents indicating meteorite contamination. It is clast rich and somewhat akin to the Apollo 14 breccias.




Figure 4: Pyroxene and olivine compositions in matrix of 72275 (from Stoeser et al. 1974 and Ryder et al. 1975).

This sample has a breccia-in-breccia texture where clasts of darker microbreccia are included in the light feldspathic matrix (figure 4). The major mineral in the matrix is feldspar. The darker areas have a higher percentage of fine matrix to clasts (Willis 1985). Portions of the light friable matrix of sample 72275 are very porous (5 to 30 percent).

The first slab cut from 72275 was the object of study by the “Consortium Indomitable” (John Wood, leader). Two additional slabs were cut from 72275 for the “breccia-pull-apart” study led by Larry Taylor (Salpas et al. 1985). Ryder (1993) provided a comprehensive review of all aspects of 72275. Recently, Nemchin et al. (2008) dated 18 zircons from 72275 ranging in age from 4.24 to 4.37 b.y., while another zircon from this boulder was found to be as old as 4.417 b.y. (Nemchin et al. 2009).

Clast Population 72275 (from Stoeser et al. 1974)

Granulitic ANT breccias	48 %
Granulitic polygonal anorthosite	3.5
Crushed anorthosite	5.1
Devitrified glass	7.9
Glass shards	0.4
Ultramafic particles	1.6
Basaltic troctolite	2.0
Pigeonite basalt	5.1
Other basaltic	2
Granite	1.6
Norite	0.4
Plagioclase	15
Mafic minerals	5.5
Opaques	1.2



Figure 5: Front and back of first slab (.42) cut from 72275. NASA# S73-32623-32624. Cube is 1 inch.

clast #1
“Marble
Cake”




Figure 6: East and West face of slab 72275,328 prepared for Taylor consortium, 1984. NASA # S84-45542 and S85-29430. Cube is 1 inch. The “Marble Cake” clast studied by the Consortium Indomitable is seen in this slab as well.




Figure 7: Thin section photomicrograph and pyroxene quadrilateral for Marble Cake clast (from Consortium Indomitabile vol 2).




Figure 8: Pyroxene and olivine composition of small granitic feldspathic clasts in 72275 (from Salpas et al. 1988).

Petrography

72275 is a polymict breccia with about 60% light porous matrix and 40% clasts. The majority of the clasts are dark aphanitic microbreccia, but also include non-mare basalt and feldspathic, plutonic fragments of the lunar crust. The light matrix of 72275 is a porous aggregate of angular mineral and lithic fragments ranging in size up to 0.1 mm (Marvin 1975). Calcic

feldspar (An_{92-98}) is the dominant mineral phase. The compositions of olivine and pyroxene mineral fragments in the matrix are given in figure 4.

The dark grey aphanitic clasts are dense and themselves microbreccias (see clasts 2 and 3 below) and equivalent to the majority of the material in the other samples of the boulder (72215, 72235, 72255).

A light colored zone through the sample is made up almost entirely of crushed non-mare basaltic material. The non-mare basalt clasts are fine-grain pigeonite basalts (equivalent to KREEPy basalt), with about equal amounts of pyroxene and plagioclase (figure 10).

There are a number of other rock types present in the clast population of 72275 including: ilmenite microgabbro, pink spinel troctolite basalt, granitic feldspathic clasts and at least one ferroan anorthosite (otherwise rare for Apollo 17) (Ryder et al. 1975, Salpas et al. 1988). Some of the larger clasts were analyzed and are discussed individually below.

Clast 1 Marble Cake Clast (,566)

The very prominent clast known as the "Marble Cake Clast" (seen in figures 6 and 27) is 3 cm in size with cataclastic gabbroic anorthosite and other material crudely interlayer with grey breccia and trace-element-rich, dark rim material (figure 7). The white core material is a mix of anorthositic norite, ilmenite microgabbro, granite and other small lithic fragments with various textures. These are swirled with vesicular glass as if lightly stirred in a marble cake (Ryder et al. 1975). The black rim of the Marble Cake clast (,80) has the composition of KREEP basalt (Blanchard et al. 1975). Nunes and Tatsumoto (1975) found that their split of the marble cake clast plotted well off of the U/Pb discordia line defined by the other boulder samples.

Clasts 2 and 3 Dark gray aphanitic clasts (,42)

These large (1 cm?) dark aphanitic clasts are seen in figures 5 and 6. Blanchard et al. (1975) found that clast 2 (,83) had a trace element pattern similar to KREEP, but Morgan et al. (1975) determined 3.44 ppb Ir (non pristine). Leich et al. (1975) obtained an Ar release pattern, but could not determine the age.

Clast 4 Pigeonite Basalt (,170)

Pigeonite basalts (later termed KREEPy Apollo 17 basalts) have equal amounts of pyroxene and plagioclase and are trace element rich. Compston et al. (1975) and Nunes and Tatsumoto (1975) studied



Figure 9: Thin section photomicrograph of typical KREEPy “pigeonite basalt in 72275.

the same pigeonite basalt (there is some confusion as to which clast this is). Compston et al. obtained an isochron age (figure 15).

Clast 5 Pigeonite Basalt (,91)

Blanchard et al. (1975), Morgan et al. (1975) analyzed another pristine clast of pigeonite basalt (figure 12). Ryder et al. (1977) and others have studied these non-mare basalts in detail and determined their mineralogy (figure 10). Unlike the Apollo 14 and 15 non-mare basalts, the “pigeonite basalts” in 72275 do not have orthopyroxene cores. Leich et al. (1975) determined




Figure 11: Pyroxene in feldspathic clast in 72275 (from Salpas et al. 1988).




Figure 10: Pyroxene and plagioclase composition in pigeonite basalt clast (KREEP). Data replotted from Ryder et al. (1975). Similar data can be found in Salpas et al. (1987).




Figure 12: Thin section photomicrograph of granu-litic feldspathic clast in 72275 (borrowed from Consortium Indomitabile). The figure illustrates calcic plagioclase surrounded by mafic minerals (see figure 8).

Table 1. Chemical composition of 72275 matrix.

reference	LSPET 73	Rose 74	Hubbard 74	Blanchard 75	Morgan 75	Jovanovic 75	Salpas 87			
weight	,2	,90	,2	,57			,413	,417	,423	
SiO ₂ %	47.54	(a) 47.31	(f)	48.3						
TiO ₂	0.91	(a) 0.94	(f)	1						
Al ₂ O ₃	17.01	(a) 16.9	(f)	16.3						
FeO	11.58	(a) 12.45	(f)	11.9			14.5	15.05	15.16	(d)
MnO	0.18	(a) 0.19	(f)	0.17						
MgO	9.35	(a) 9.47	(f)	10.3						
CaO	11.71	(a) 11.72	(f)	11			10.1	10.3	12.1	(d)
Na ₂ O	0.38	(a) 0.35	(f) 0.36	(c) 0.44			0.42	0.38	0.37	(d)
K ₂ O	0.28	(a) 0.22	(f) 0.28	(b) 0.25						
P ₂ O ₅	0.35	(a) 0.38	(f)							
S %	0.08	(a)								
sum										
Sc ppm		40	(f)	44.7	(d)		45.7	48.6	49.8	(d)
V		75	(f)							
Cr		2330	(f)	2395	(d)		3062	3088	3255	(d)
Co		37	(f)	30.4	(d)		31.3	33.3	35.3	(d)
Ni	67	(a) 127	(f)	75	(d)	95	12	55	<110	(d)
Cu		5.4	(f)							
Zn	3	(a)				2.7				
Ga		3.2	(f)							
Ge ppb					406	(e)				
As										
Se ppb					34	(e)				
Rb	8.7	(a) 4.6	(f) 8.97	(b)	5.9	(e)	13	12	14	(d)
Sr	121	(a) 135	(f) 123	(b)			138	93	<160	(d)
Y	129	(a) 88	(f)							
Zr	613	(a) 545	(f) 605	(b)			600	765	700	(d)
Nb	32	(a) 24	(f)							
Mo										
Ru										
Rh										
Pd ppb										
Ag ppb					0.74	(e)				
Cd ppb					13	(e)				
In ppb										
Sn ppb										
Sb ppb					1.17	(e)				
Te ppb					4.14	(e)				
Cs ppm					0.255	(e)	0.37	0.4	0.44	(d)
Ba		330	(f) 350	(b)			370	400	400	(d)
La		35	(f) 41	(b) 50.5	(d)		47.9	50.2	52.3	(d)
Ce			106	(b) 130	(d)		129	133	139	(d)
Pr										
Nd			67.4	(b)			80	81	85	(d)
Sm			18.8	(b) 24.6	(d)		22.2	23.5	25.5	(d)
Eu			1.49	(b) 1.57	(d)		1.62	1.66	1.68	(d)
Gd			23.4	(b)						
Tb				4.9	(d)		4.59	4.97	5.1	(d)
Dy			23.2	(b)						
Ho										
Er			13.7	(b)						
Tm										
Yb		9.2	(f) 11.6	(b) 15	(d)		13.5	13.9	13.1	(d)
Lu			1.71	(b) 2.01	(d)		1.73	1.8	1.9	(d)
Hf			14.6	(b) 16.5	(d)		16.4	17.2	17.9	(d)
Ta				1.7	(d)		1.55	1.66	1.58	(d)
W ppb										
Re ppb					0.225	(e)				
Os ppb										
Ir ppb					2.26	(e)	<2	<2	<2	(d)
Pt ppb										
Au ppb					0.82	(e)	<5	<7	<6	(d)
Th ppm			5.29	(b)			5.52	5.46	6.01	(d)
U ppm			1.56	(b)	1.5	(e) 1.6	1.3	1.58	1.26	(d)

technique: (a) XRF, (b) ID/MS, (c) AA, (d) INAA, (e) RNAA, (f) varied

Table 2. Chemical composition of clasts in 72275.

	anorthosite		pigeonite basalt		marble cake			KREEP basalt			granulite	
reference	Salpas 88		Blanchard 75	Morgan 75	Blanchard 75			Salpas 87			Salpas 88	
weight	FAN ,350		PB ,91	PB ,91	clast,80	clast 1		,385	,357	,427b	,397	,433
SiO2 %			48	(a)	47	47	47		51.3	48.3		
TiO2			1.4	(a)	1.8	1.1	1.8		1.54	1.2	0.22	0.15
Al2O3			13.5	(a)	17.9	18.2	23.5		14.5	12.5	26.2	24.6
FeO	0.485	(a)	15	(a)	10.3	10.9	7.4	15.18	13.9	16.5	5.71	5.1
MnO			0.156	(a)	0.104	0.17	0.08		0.17	0.22		
MgO			10	(a)	9.43	9.14	5.24		6.8	11.4	7.9	8
CaO	19.2	(a)	10.5	(a)	11.7	11.2	14.2	9.1	10.8	9.5	14.8	14.2
Na2O	0.456	(a)	0.29	(a)	0.39	0.63	0.36	0.35	0.51	0.415	0.353	0.362
K2O			0.25	(a)	0.47	0.49	0.32					
P2O5												
S %												
sum												
Sc ppm	1.12	(a)	61	(a)	34	26.3	25	50	51	45.5	7.81	8.24
V									97	135	20	24
Cr	46.6	(a)	3147	(a)	3147			3170	1960	4420	842	881
Co	0.44	(a)	37	(a)	28	22.5	18.7	35.1	30.9	46.4	39.3	30.6
Ni	<7	(a)		43	(b)	130		50	<80	112	455	422
Cu												
Zn				2.7	(b)							
Ga												
Ge ppb				1290	(b)							
As												
Se				0.23	(b)							
Rb				8	(b)			14	12	12		
Sr	205	(a)						93	92	98	160	160
Y												
Zr								800	610	540		
Nb												
Mo												
Ru												
Rh												
Pd ppb												
Ag ppb				0.58	(b)							
Cd ppb				8.3	(b)							
In ppb												
Sn ppb												
Sb ppb				2.87	(b)							
Te ppb				7.8	(b)							
Cs ppm	0.016	(a)		0.355	(b)			0.55	0.4	0.3	0.19	0.23
Ba	40	(a)						440	500	365	72	87
La	0.567	(a)	48	(a)	78	78	48	52.5	61.7	46.2	3.66	4.72
Ce	1.48	(a)	131	(a)	213	206	131	140	155	121	10.1	12.6
Pr												
Nd	<2.5	(a)						92	108	75	5.7	6.2
Sm	0.228	(a)	23	(a)	36	36	22.5	23.8	28.9	22.3	1.56	1.93
Eu	0.928	(a)	1.58	(a)	2.14	2.1	1.81	1.62	1.87	1.45	0.835	0.86
Gd												
Tb	0.045	(a)	4.5	(a)	7.7	7.7	4.7	4.9	5.82	4.31	0.375	0.49
Dy												
Ho												
Er												
Tm												
Yb	0.125	(a)	11.9	(a)	24	25.4	13.9	13.8	15.5	12.4	1.69	2.06
Lu	0.02	(a)	1.75	(a)	3.5	3.5	2.04	1.83	2.18	1.67	0.238	0.292
Hf	0.133	(a)	18	(a)	19.8	25.1	14	17.4	20.5	15.9	1.46	1.98
Ta	0.015	(a)				3.5		1.62	1.9	1.37	0.302	0.309
W ppb												
Re ppb				0.007	(b)							
Os ppb												
Ir ppb	<2	(a)		0.023	(b)			<2			16.4	14
Pt ppb												
Au ppb	<0.8	(a)		0.045	(b)			<7			6.8	6.5
Th ppm	0.047	(a)				12.8		5.98	6.73	5.25	1.17	2.06
U ppm	0.02	(a)		1.5	(b)			1.3	1.95	1.45	0.34	0.37
technique	(a) INAA, (b) RNAA											

an Ar release plateau, but were not able to determine an age.

KREEP basalt clasts

Ryder (1977) and Salpas et al. (1987) describe several KREEP basalt clasts, several of which had a subophitic basaltic texture (figure 9). They equate these to the KREEPy pigeonite basalt clasts previously studied. A large number of these clasts, including two with pristine igneous texture, were analyzed by Salpas et al. (1987). Shih et al. (1992) were able to obtain one of the brecciated KREEP basalt clasts and date it by both Rb-Sr and Sm-Nd (figures 16 and 17).

Ferroan Anorthosite Clast (,350)

A unique fragment of pristine ferroan anorthosite was studied by Salpas et al. (1988). It was 3 x 4 x 5 mm and composed of 95% plagioclase (An_{96}) and 5% pyroxene (figure 12). The chemical composition is given in table 2 and this clast is apparently plutonic and pristine ($Ir < 2$ ppb). Since this is what we think the lunar crust was made of, we are surprised to not find more fragments of this kind in the ejecta of large impacts.

Granulitic ANT Clasts

The Consortium Indomitabile coined the acronym ANT for the small fragments of plagioclase-rich, potash and phosphorous-poor rocks whose mineralogies vary over the range anorthosite-norite-troctolite (*never mind the otherwise small grain size*). In 72275 the textures of these feldspathic clasts are that of an annealed granulite (figure 12). Salpas et al. (1988) discovered several that were large enough to analyze (table 2, figure 13). However, they were all high in Ir.

Granite Clast

Meyer et al. (1996) featured a large coarse-grained granite clast on the cover of meteoritics (figure 30).

Mineralogy

The mineral fragments in the matrix are, in order of abundance: plagioclase, olivine, orthopyroxene, pigeonite, Ca-rich clinopyroxene, ilmenite, spinel, cristobalite, barium K-feldspar, Fe metal, troilite, zircon and armalcolite (Stoeser et al. 1974).

Chemistry

Chemical analyses of the matrix of 72275 are given in table 1 and of selected clasts in table 2. In addition there are a number of analyses of small clasts in thin




Figure 13: Normalized rare-earth-element diagram for matrix and selected clasts in breccia sample 72275. Data for matrix and pigeonite basalt is from Blanchard et al. (1975), and for clasts Salpas et al. (1987, 1988).

section by broad-beam, electron probe microanalyses in the papers by Ryder et al. (1975) and Stoeser et al. (1975). Salpas et al. (1987, 1988) present the analyses of a large number of clasts (mostly KREEP basalt and granulitic feldspathic clasts (ranges shown in figure 13).

The composition of the light matrix of 72275 is broadly similar to that of the pigeonite basalt clasts (Salpas et al. 1987). There is a hint of slightly elevated Al, which would be expected by the presence of anorthositic clasts.

The trace siderophile element ratios (Morgan et al. 1975) are not exactly what is expected for Serenitatis ejecta. Ge is very high, and there is evidence that this is from the high KREEP component (which is itself surprisingly high in Ge).




Figure 14: Ar-Ar release pattern for 72275,80. Plateau age is 3.99 ± 0.03 b.y. (from Leich et al. 1975).

Radiogenic age dating

Compston et al. (1975) and Shih et al. (1992) found that the age of the KREEPy basalt clasts in 72275 to be about 200 m.y. older than the ages of KREEP basalts from Apollo 15 (such as 15382, 15386). Nunes and Tatsumoto (1975) attempted to date various chips of 72275 for the consortium by U-Th-Pb, but learned instead that the breccia and its clasts has suffered extensive movement of Pb. Leich et al. (1975) also attempted to obtain Ar-Ar dates, but found that plateau ages were ill defined except for one case. (please note that dates reported here are the original data and require correction for new decay constants)

Summary of Age Data for 72275

	Ar-Ar	Rb-Sr
Leich et al. 1975	3.99 ± 0.04 b.y.	
Compston et al. 1975		4.01 ± 0.04
Shih et al. 1992		4.13 ± 0.08
Nemchin et al. 2008	$4.24 - 4.37$ b.y.	
Grange et al. 2009		

Caution: Beware decay constants!




Figure 15: Rb-Sr isochron diagram of pigeonite clast in 72275 (from Compston et al. 1975).




Figure 16: Rb-Sr isochron diagram for KREEP basalt clast in 72275 (from Shih et al. 1992).





Figure 17: Sm-Nd isochron diagram for KREEP basalt clast in 72275 (from Shih et al. 1992).

Sm-Nd

4.08 ± 0.07	Rim of Marble Cake Pigeonite basalt, 171 KREEP basalt clast B-1 Pb/Pb Zircons
-----------------	--



Leich et al. (1975) determined the exposure age of 72275 and other samples of the same boulder by a number of techniques, of which ^{81}Kr are the most reliable (figure 18). The conclusion is that 72275 (from the top of the boulder) has been exposed to cosmic rays for 52.5 ± 1.4 m.y.

72275

,102

,337

,328

W₁Si

Figure 19: Sketch of slabs cut from 72275, 102 in 1984

Pearce et al. (1974), Brecher et al. (1974) and Banerjee and Swits (1975) determined the magnetic properties of 72275. Housley et al. (1977) included 72275 in their study of ferromagnetic resonance of lunar samples (it had none).

Goswami and Hutcheon (1975) and Goswami et al. (1977) studied the track densities in minerals from 72275. Charette and Adams (1977) have reported the spectra.

Figure 19: Sketch of slabs cut from 72275,102 in 1984.







Figure 21: Photo and map of west face of second slab (,328) of 72275 (Willis 1985). S84-41208. The large clast is the other half of the Marble Cake Clast studied by Marvin et al. 1973. (see figure27)



72275,328
East Face





Figure 22: Photo and map of east face of second slab (,328) of 72275 (Willis 1985). S84-39544.



West Face



Figure 23: Photo and map of west side of third slab (, 337) of 72275 (Willis 1985). S84-39547




Figure 24: Photo and map of east side of third slab (, 337) of 72275 (Willis 1985). S84-39546.



72275,102
West Face




Figure 25: Photo and map of west face of 72275,102 (end piece). S84-39545. From Salpas et al. 1985.





Figure 26: Photo of slab (,337) after subdivision. Cube is 1 inch. S84-46140. (see figure 24)



Marble Cake Clast

Figure 27: Photo of clasts extracted from slab 72275,328. S97-16861. Cube is 1 cm. (see figure 21). Subsample ,566 is the other half of the Marble Cake Clast studied by the Wood consortium (Ryder 1992).




Figure 28: Thin section photomicrograph of clast 72275,493. S84-46669. Scale unknown.




Figure 29: Thin section photo of clast 72275,497(basalt). S84-4667. Scale unknown.



Figure 30: False color image by BSE of granite clast in 72275 (Meyer et al. 1996). Field of view is 4 mm. Red is K-feldspar, yellow is plagioclase and black is silica.

References for 72275

Adams J.B. and Charette M.P. (1975) Spectral reflectance of highland rock types at Apollo 17: Evidence from Boulder 1, Station 2. *The Moon* **14**, 483-489.

Arvidson R., Drozd R., Guinness E., Hohenberg C., Morgan C., Morrison R. and Oberbeck V. (1976) Cosmic ray exposure ages of Apollo 17 samples and the age of Tyco. *Proc. 7th Lunar Sci. Conf.* 2817-2832.

Banerjee S.K. and Swits G. (1975) Natural remanent magnetization studies of a layered breccia boulder from the lunar highland region. *The Moon* **14**, 473-481.

Processing

72275 is friable and broke into pieces on the return to Earth (figure 1). A partial slab (,42) was cut to expose the prominent clast for the Consortium Indomitable in 1974. This saw cut was completed and two additional slabs were cut for the Larry Taylor “breccia-pull-apart” project in 1984. A partial “flow diagram” of processing of 72275 is provided. A piece of 72275 is on public display at the Langley Research Center in Virginia.


There are 110 thin sections of 72275. Some are used in the educational thin section packages; many are of the clasts that have been extracted from the slabs.

Banerjee S.K., Hoffman K. and Swits G. (1974a) Remanent magnetization directions in a layered boulder from the South Massif. *Proc. 5th Lunar Sci. Conf.* 2873-2881.

Banerjee S.K., Hoffman K. and Swits G. (1974b) Reversed polarity remanent magnetization in a layered boulder near South Massif (abs). *LS V*, 32-34.

Blanchard D.P., Haskin L.A., Jacobs J.W., Brannon J.C. and Korotev R.L. (1975) Major and trace element chemistry of Boulder 1 at Station 2, Apollo 17. *The Moon* **14**, 359-371.

Compston W., Foster J.J. and Gray C.M. (1975) Rb-Sr ages of clasts from within Boulder 1, Station 2, Apollo 17. *The Moon* **14**, 445-462.



Dalrymple G.B. and Ryder G. (1996) Argon-40/argon-39 age spectra of Apollo 17 highlands breccia samples by laser step heating and the age of the Serenitatis basin. *J. Geophys. Res.* 101, 26069-26084.

Fruchter J.S., Rancitelli L.A. and Perkins R.W. (1975) Primordial radionuclide variations in the Apollo 15 and 17 deep core samples and in Apollo 17 igneous rocks and breccias. *Proc. 6th Lunar Sci. Conf.* 1399-1406.

Goswami J.N. and Hutcheon I.D. (1975) Cosmic ray exposure history and compaction age of Boulder 1 from Station 2. *The Moon* 14, 395-405.

Goswami J.N., Braddy D. and Price P.B. (1976a) Microstratigraphy of the lunar regolith and compaction ages of lunar breccias. *Proc. 7th Lunar Sci. Conf.* 55-74.

Goswami J.N., Braddy D. and Price P.B. (1976b) Microstratigraphy of the lunar regolith and compaction ages of lunar breccias (abs). *LS VII*, 328-330.

Grange M., Nemchin A.A., Pidgeon R.T. and Meyer C. (2009) Early history of the moon: Zircon perspective (abs#1473). *Lunar Planet. Sci. XL*, Luanr Planetary Institute, The Woodlands.

Higuchi H. and Morgan J.W. (1975) Ancient meteoritic component in Apollo 17 boulders. *Proc. 6th Lunar Sci. Conf.* 1625-1651.

Hubbard N.J., Rhodes J.M., Wiesmann H., C.-Y. Shih and B.M. Bansal (1974) The chemical definition and interpretation of rock types returned from the non-mare regions of the moon. *Proc. 5th Lunar Sci. Conf.* 1227-1246.

Hutcheon I.D., MacDougall D. and Stevenson J. (1974b) Apollo 17 particle track studies: surface residence times and fission track ages for orange glass and large boulders. *Proc. 5th Lunar Sci. Conf.* 2597-2608.

James O.B. (1982) Subdivision of the Mg-suite plutonic rocks into Mg-norites and Mg-gabbro-norites (abs). *LPS XIII*, 360-362.

James O.B. and Flohr M.K. (1983) Subdivision of the Mg-suite noritic rocks into Mg-gabbro-norites and Mg-norites. *Proc. 13th Lunar Planet. Sci. Conf.*, *J. Geophys. Res.*, A603-A614.

Jovanovic S. and Reed G.W. (1975a) Cl and P₂O₅ systematics: Clues to early lunar magmas. *Proc. 6th Lunar Sci. Conf.* 1737-1751.

- Jovanovic S. and Reed G.W. (1975b) Soil breccia relationships and vapor deposits on the moon. *Proc. 6th Lunar Sci. Conf.* 1753-1759.
- Jovanovic S. and Reed G.W. (1975c) History of Boulder 1 at Station 2, Apollo 17 based on trace element interrelationships. *The Moon* 14, 385-393.
- Jovanovic S. and Reed G.W. (1975d) Studies on regolith processes: Apollo 15 and 17 labile trace element implications (abs). *Lunar Sci.* VI, 451-453.
- Jovanovic S. and Reed G.W. (1980a) Candidate samples for the earliest lunar crust. *Proc. Conf. Lunar Highlands Crust, Geochim. Cosmochim. Acta, Suppl. 12.* Pergamon Press. 101-111.
- Keith J.E., Clark R.S. and Bennett L.J. (1974a) Determination of natural and cosmic ray induced radionuclides in Apollo 17 lunar samples. *Proc. 5th Lunar Sci. Conf.* 2121-2138.
- Keith J.E., Clark R.S. and Bennett L.J. (1974b) Determination of natural and cosmic ray induced radionuclides in Apollo 17 lunar samples (abs). *LS V*, 402-404.
- Knoll H.-D. and Stöffler D. (1979) Characterization of the basic types of lunar highland breccias by quantitative textural analysis (abs). *LPS X*, 673-675.
- Leich D.A., Kahl S.B., Kirschbaum A.R., Niemeyer S. and Phinney D. (1975a) Rare gas constraints on the history of Boulder 1, Station 2, Apollo 17. *The Moon* 14, 407-444.
- MacDougall D., Rajan R.S., Hutcheon I.D. and Price P.B. (1973) Irradiation history and accretionary processes in lunar and meteoritic breccias. *Proc. 4th Lunar Sci. Conf.* 2319-2336.
- MacDougall D., Hutcheon I.D. and Price P.B. (1974) Irradiation records in orange glass and two boulders from Apollo 17 (abs). *Lunar Sci.* V, 483-485.
- Marvin U.B. (1971) Lunar niobian rutile. *Earth Planet. Sci. Lett.* 11, 7-9.
- Marvin U.B. (1975) The Boulder. *The Moon* 14, 315-326.
- Meyer C., Williams I.S. and Compston W. (1996) Uranium-lead ages for lunar zircons: Evidence for a prolonged period of granophyre formation from 4.32 to 3.88 Ga. *Meteor. & Planet. Sci.* 31, 370-387.
- Morgan J.W., Ganapathy R., Higuchi H., Krahenbuhl U. and Anders E (1974a) Lunar basins: Tentative characterization of projectiles, from meteoritic elements in Apollo 17 boulders. *Proc. 5th Lunar Sci. Conf.* 1703-1736.
- Morgan J.W., Ganapathy R., Higuchi H., Krahenbuhl U. and Anders E. (1974b) Lunar basins: Tentative characterization of projectiles, from meteoritic elements in Apollo 17 boulders (abs). *Lunar Sci.* V, 526-528.
- Morgan J.W., Higuchi H. and Anders E. (1975b) Meteoritic material in a boulder from the Apollo 17 site: Implications for its origin. *The Moon* 14, 373-383.
- Mori H. and Takeda H. (1980) Thermal and deformational history of diogenites and a lunar norite, as determined by electron microscopy and crystallography (abs). *LPS XI*, 743-745.
- Mori H., Takeda H. and Miyamoto M. (1982) Comparison of orthopyroxenes in lunar norites and diogenites (abs). *LPS XIII*, 540-541.
- Nemchin A.A., Pidgeon R.T., Whitehouse M.J., Vaughan J.P. and Meyer C. (2008) SIMS study of zircons from Apollo 14 and 17 breccias: Implications for the evolution of lunar KREEP. *Geochim. Cosmochim. Acta* 72, 668-689.
- Nemchin A.A., Timms N., Pidgeon R.T., Geisler T., Reddy S. and Meyer C. (2009) Timing of crystallization of the lunar magma ocean constrained by the oldest zircon. *Nature Geoscience* doi 10.1038/NCEO419
- Nunes P.D. and Tatsumoto M. (1975a) U-Th-Pb systematics of selected samples from Apollo 17, Boulder I, Station 2. *The Moon* 14, 463-471.
- Nunes P.D., Tatsumoto M. and Unruh D.M. (1974b) U-Th-Pb systematics of some Apollo 17 lunar samples and implications for a lunar basin excavation chronology. *Proc. 5th Lunar Sci. Conf.* 1487-1514.
- Nyquist L.E. and Shih C-Y. (2008) Chronology and provenance of lunar KREEP: A 4.0 or 4.1 Ga age for Serenitatis? (abs) Early Solar System Impact Bombardment Workshop. Lunar Planet. Institute, Houston.
- Palme H., Baddenhausen H., Blum K., Cendales M., Dreibus G., Hofmeister H., Kmse H., Palme C., Spettel B. Vilesek E. and Wanke H. (1978) New data on lunar samples and achondrites and a comparison of the least fractionated samples from the earth, the moon, and the eucrite parent body. *Proc. 9th Lunar Planet. Sci. Conf.* 25-57.
- Reed V.S. and Wolfe E.W. (1975) Origin of the Taurus-Littrow Massifs. *Proc. 6th Lunar Sci. Conf.* 2443-2462.
- Ryder G. (1983) Nickel in olivines and parent magmas of lunar pristine rocks. *In* Workshop on Pristine Highlands

Rocks and the Early History of the Moon (Longhi and Ryder, eds.) LPI Tech Rept. 83-02. The Lunar and Planetary Institute, Houston, 66-68.

Ryder G. (1984) Most olivine in the lunar highlands is of shallow origin (abs). LPS XV, 707-708.

Ryder G. (1992) A distinct poikilitic impact melt rock from the Apollo 17 landing site that is not from the Serenitatis melt sheet. (abs) Meteoritics 27, 284.

Ryder G. (1993a) The Apollo 17 samples: The massifs and landslide. In Workshop on Geology of the Apollo 17 Landing Site. LPI Tech. Rpt. 92-09, 48-49.

Ryder G. (1993b) Impact melt breccias at the Apollo 17 landing site. In Workshop on Geology of the Apollo 17 Landing Site. LPI Tech. Rpt. 92-09, 49-50.

Ryder G. (1993c) **Catalog of Apollo 17 rocks: Stations 2 and 3.** Curators Office JSC#26088.

Ryder G., Stoesser D.B., Marvin U.B. and Bower J.F. (1975a) Lunar granites with unique ternary feldspars. Proc. 6th Lunar Sci. Conf. 435-449.

Ryder G., Stoesser D.B., Marvin U.B., Bower J.F. and Wood J.A. (1975b) Boulder 1, Station 2, Apollo 17: Petrology and petrogenesis. The Moon 14, 327-357.

Ryder G. and Norman M.D. (1979a) Catalog of pristine non-mare materials Part 1. Non-anorthosites, revised. NASA-JSC Curatorial Facility Publ. JSC 14565, Houston. 147 pp. 72255

Ryder G. and Spudis P. (1980) Volcanic rocks in the lunar highlands. Proc. Conf. Lunar Highlands Crust, 353-375. GCA 12, Lunar Planetary Inst.

Ryder G., Norman M.D. and Score R.A. (1980a) The distinction of pristine from meteorite-contaminated highlands rocks using metal compositions. Proc. 11th Lunar Planet. Sci. Conf. 471-479.

Ryder G., Norman M.D. and Score R.A. (1980b) Ni, Co content of metal grains for the identification of indigenous rocks (abs). LPS XI, 968-970.

Ryder G., Norman M.D. and Taylor G.J. (1997) The complex stratigraphy of the highland crust in the Serenitatis region of the Moon inferred from mineral fragment chemistry. *Geochim. Cosmochim. Acta* 61, 1083-1105.

Salpas P.A. and Taylor L.A. (1985) Basalt clasts in breccia 72275: Examples of pre-mare volcanism (abs). Lunar Planet. Sci. XVI, 728-729. Lunar Planetary Institute, Houston.

Salpas P.A., Willis K.J. and Taylor L.A. (1985) Breccia Guidebook No. 8, 72275. Curatorial Branch Publication 71, JSC 20416 pp. 43.

Salpas P.A., Taylor L.A. and Lindstrom M.M. (1986a) Apollo 17 KREEPy basalts: Pristine basaltic breccias (abs). LPS XVII, 748-749.

Salpas P.A., Taylor L.A. and Lindstrom M.M. (1986b) The first Apollo 17 ferroan anorthosite: Its significance relative to Mg-suite highland clasts (abs). LPS XVII, 752-753.

Salpas P.A., Lindstrom M.M. and Taylor L.A. (1987) Highland materials at Apollo 17: Contributions from 72275. *Proc. Lunar and Planet. Sci. Conf.* 18th, 11-19.

Salpas P.A., Taylor L.A. and Lindstrom M.M. (1987) Apollo 17 KREEPy basalts: Evidence for Nonuniformity of KREEP. *Proc. Lunar Planet. Sci. Conf.* 17th, E340-E348.

Salpas P.A., Lindstrom M.M. and Taylor L.A. (1988) Highland materials at Apollo 17: Contributions from 72275. *Proc. Lunar Planet. Sci. Conf.* 18th, 11-19.

Schaeffer O.A., Warasila R. and Labotka T.C. (1982a) Ages of Serenitatis breccias: Lunar breccias and soils and their meteoritic analogs. LPI Tech. Rept. 82-02, 123-125.

Schaeffer O.A., Warasila R. and Labotka T.C. (1982b) Ages of Serenitatis breccias (abs). LPS XIII, 685-686.

Schmitt H.H. (1975) Geological model for Boulder 1 at Station 2, South Massif, Valley of Taurus-Littrow. The Moon 14, 491-504.

Shih C.-Y., Bansal B.M., Wiesmann H. and Nyquist L.E. (1990a) Rb-Sr and Sm-Nd isotopic studies of an Apollo 17 KREEPy basalt (abs). LPS XXI, 1148-1149.

Shih C.-Y., Nyquist L.E., Bansal B.M. and Wiesmann H. (1992) Rb-Sr and Sm-Nd chronology of an Apollo 17 KREEP basalt. *Earth Planet. Sci. Lett.* 108, 203-215.

Simonds C.H., Phinney W.C. and Warner J.L. (1974) Petrography and classification of Apollo 17 non-mare rocks with emphasis on samples from the Station 6 boulder. Proc. 5th Lunar Sci. Conf. 337-353.

Stoesser D.B., Marvin U.B., Wood J.A., Wolfe R.W. and Bower J.F. (1974a) Petrology of a stratified boulder from South Massif, Taurus-Littrow. Proc. 5th Lunar Sci. Conf. 355-377.

Stoeser D.B., Wolfe R.W., Marvin U.B., Wood J.A. and Bower J.F. (1974b) Petrographic studies of a boulder from the South Massif (abs). LS V, 743-745.

Stoeser D.B., Wolfe R.W., Wood J.A. and Bower J.F. (1974c) Petrology and petrogenesis of boulder 1. *In* Interdisciplinary Studies of Samples from Boulder 1, Station 2, Apollo 17. Volume 1, Consortium Indomitable. Smithsonian Astrophysical Observatory. *Also* Lunar Science Institute Cont. no. 210D, 35-109.

Stoeser D.B., Marvin U.B. and Bower J.F. (1974d) Petrology and petrogenesis of boulder 1. *In* Interdisciplinary Studies of Samples from Boulder 1, Station 2, Apollo 17. Volume 2, Consortium Indomitable. Smithsonian Astrophysical Observatory. *Also* Lunar Science Institute Cont. no. 21 ID, 1-59.
72215 72235 72275

Stoeser D.B., Ryder G. and Marvin U.B. (1975) Lunar granite clasts with unique ternary feldspars (abs). LS VI, 780-782.

Spudis P.D. and Ryder G. (1981) Apollo 17 impact melts and their relation to the Serenitatis basin. *In* Proc. of the Conf. on Multi-Ring Basins. Proc. Lunar Planet. Sci. 12A - Geochim. Cosmochim. Acta, Suppl. 15. Pergamon Press. 133-148.

Takeda H. and Ishii T. (1975) Typical processes of exsolution, decomposition and inversion of pyroxenes and its bearing on thermal history of lunar rocks (abs). LS VI, 795-797.

Takeda H. and Miyamoto M. (1976) Characterization of crust formation on a parent body of achondrites and the Moon by pyroxene crystallography and chemistry (abs). LS VII, 846-848.

Takeda H., Miyamoto M., Ishii T. and Reid A.M. (1976) Characterization of crust formation on a parent body of achondrites and the Moon by pyroxene crystallography and chemistry. Proc. 7th Lunar Sci. Conf., 3535-3548.

Takeda H., Mori H. and Miyamoto M. (1982) Comparison of thermal history of orthopyroxenes between lunar norites 78236, 72255, and diogenites. Proc. 13th Lunar Planet. Sci. Conf. A124-A130.

Willis K.J. (1985) Three lithologic units of 72275 (abs). Lunar Planet. Sci. XVI, 910-911. Lunar Planetary Institute, Houston.

Willis K.J. (1985) Mapping of five new faces of 72275. Lunar sample Newsletter #42, Append. 1, 9-23. Curator's Office, JSC.

Winzer S.R., Nava D.F., Schuhmann S., Lum R.K.L. and Philpotts J.A. (1975a) Origin of the Station 7 boulder: A note. Proc. 6th Lunar Sci. Conf. 707-710.

Wolf R., Woodrow A. and Anders E. (1979) Lunar basalts and pristine highland rocks: Comparison of siderophile and volatile elements. Proc. 10th Lunar Planet. Sci. Conf. 2107-2130.

Wolfe E.W. et al. (1981) The geologic investigation of the Taurus-Littrow valley: Apollo 17 landing site. US Geol. Survey Prof. Paper 1080.